

ON ENTROPY, REGULARITY AND RIGIDITY FOR CONVEX REPRESENTATIONS OF HYPERBOLIC MANIFOLDS

A. SAMBARINO

ABSTRACT. Given a convex representation $\rho : \Gamma \rightarrow \mathrm{PGL}(d, \mathbb{R})$ of a convex co-compact group Γ of \mathbb{H}^k we find upper bounds for the quantity αh_ρ , where h_ρ is the entropy of ρ and α is the Hölder exponent of the equivariant map $\partial_\infty \Gamma \rightarrow \mathbb{P}(\mathbb{R}^d)$. We also give rigidity statements when the upper bound is attained. We then study Hitchin representations and prove that if $\rho : \pi_1 \Sigma \rightarrow \mathrm{PSL}(d, \mathbb{R})$ is in the Hitchin component then $\alpha h_\rho \leq 2/(d-1)$ (where α is the Hölder exponent of the map $\zeta : \partial_\infty \mathbb{H}^2 \rightarrow \mathcal{P}$) with equality if and only if ρ is Fuchsian.

CONTENTS

1. Introduction	1
2. Reparametrizations and Thermodynamic Formalism	10
3. $\mathrm{CAT}(-1)$ spaces	12
4. Convex representations	14
5. Proof of Theorem A	17
6. Cross ratios: General Lemmas and a Theorem of Labourie	20
7. The cross ratio of a Hölder cocycle	21
8. Convex representations and cross ratios	23
9. Proximal Representations of semi-simple Lie groups	24
10. Computing the rank of $\mathbb{B}_{\mathbb{H}}^\alpha$	24
11. Proofs of Theorem B, Corollary 1.5 and Corollary 1.6	27
12. Hyperconvex representations and Theorem C	29
13. Proofs of Theorem D and Corollary 1.12	31
14. Proof of Corollary 1.13	32
References	34

1. INTRODUCTION

Consider a $\mathrm{CAT}(-1)$ space X . Its visual boundary $\partial_\infty X$ is equipped with a natural metric, called a visual metric. This metric depends on the choice of a point in X , different points induce bi-Lipschitz equivalent metrics.

Consider now a convex co-compact action of a hyperbolic group Γ on X . An important invariant for this action is the Hausdorff dimension h_Γ for a visual metric, of the limit set L_Γ of Γ on the visual boundary $\partial_\infty X$ of X .

Several rigidity statements have been found concerning lower bounds on this Hausdorff dimension. For example, Bourdon [7] proved that if $\Gamma = \pi_1 M$ where M

The author was partially supported by the European Research Council under the *European Community's* seventh Framework Program (FP7/2007-2013)/ERC grant agreement n FP7-246918.

is a closed k -dimensional manifold modeled on \mathbb{H}^k , then $h_\Gamma \geq k - 1$ with equality only if the action of Γ on X preserves a totally geodesic copy of \mathbb{H}^k . We refer the reader to Courtois [11] for a more detailed exposition on this problem.

Given two convex-co-compact actions $\rho_i : \Gamma \rightarrow \text{Isom } X_i$ $i = 1, 2$ on $\text{CAT}(-1)$ spaces X_i , there is an obvious relation between the Hausdorff dimensions of their limit sets. Let $\xi : L_{\rho_1\Gamma} \rightarrow L_{\rho_2\Gamma}$ be the Hölder-continuous equivariant mapping. From the definition of Hausdorff dimension one obtains

$$\alpha h_{\rho_2} \leq h_{\rho_1} \quad (1)$$

where α is the Hölder exponent of ξ , i.e. $d(\xi(x), \xi(y)) \leq Kd(x, y)^\alpha$ for some $K > 0$. In this case we will say that ξ is α -Hölder.

The main purpose of this work is to extend inequality (1) for convex representations $\Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ and give rigidity results when the equality holds. In order to do so we will exploit the well known fact that h_Γ is also a dynamical invariant.

Consider the geodesic flow of $\Gamma \backslash X$,

$$\phi = \{\phi_t : \Gamma \backslash UX \rightarrow \Gamma \backslash UX\}_{t \in \mathbb{R}}.$$

The topological entropy of ϕ coincides with the Hausdorff dimension h_Γ (Sullivan [22], see also Bourdon [6]). The fact that Γ is convex-co-compact is equivalent to the fact the non wandering set of ϕ , denoted from now on $U\Gamma$, is compact. Moreover, $\phi|_{U\Gamma}$ has very nice dynamical properties coming from the negative curvature of X , namely it is a topological Anosov flow (see Definition 2.4). Its topological entropy can then be computed by counting how many periodic orbits the flow $\phi|_{U\Gamma}$ has: for non torsion $\gamma \in \Gamma$ denote by

$$|\gamma| = \inf_{p \in X} d_X(p, \gamma p)$$

the length of the closed geodesic of $\Gamma \backslash X$ determined by the conjugacy class $[\gamma]$ of γ , then

$$h_\Gamma = \lim_{t \rightarrow \infty} \frac{\log \#\{[\gamma] \in [\Gamma] : |\gamma| \leq t\}}{t}.$$

Our object of study is the following:

Definition 1.1. We will say that a representation $\rho : \Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ is *convex* if there exist two ρ -equivariant Hölder-continuous maps

$$\xi : \partial_\infty \Gamma \rightarrow \mathbb{P}(\mathbb{R}^d) \text{ and } \xi^* : \partial_\infty \Gamma \rightarrow \mathbb{P}(\mathbb{R}^{d^*})$$

such that $\xi(x) \oplus \ker \xi^*(y) = \mathbb{R}^d$ whenever $x, y \in \partial_\infty \Gamma$ are distinct.

We will study 2 different entropies one can define for a convex representation. For $g \in \text{PGL}(d, \mathbb{R})$ denote by $\lambda_1(g)$ the logarithm of the spectral radius of g . The *spectral entropy* of a convex representation $\rho : \Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ is defined by

$$h_\rho = \lim_{t \rightarrow \infty} \frac{1}{t} \log \#\{[\gamma] \in [\Gamma] : \lambda_1(\rho\gamma) \leq t\},$$

and the *Hilbert entropy* of ρ is defined by

$$H_\rho = \lim_{t \rightarrow \infty} \frac{1}{t} \log \#\left\{[\gamma] \in [\Gamma] : \frac{\lambda_1(\rho\gamma) - \lambda_d(\rho\gamma)}{2} \leq t\right\}$$

where $\lambda_d(\rho\gamma)$ is the log of the modulus of the smallest eigenvalue of $\rho\gamma$.

One has the following Proposition:

Proposition 1.2 ([20], see also Bridgeman-Canary-Labourie-S. [10]). *The spectral entropy of an irreducible convex representation of a (finitely generated non elementary) hyperbolic group is finite and positive.*

If V is a finite dimensional vector space we will consider the distance $d_{\mathbb{P}}$ on $\mathbb{P}(V)$ coming from an euclidean distance on V . An important remark is that the entropy h_{ρ} of a convex representation is not necessarily the Hausdorff dimension of $\xi(\partial_{\infty}\Gamma)$ (see Remark 1.4 below). Our first result is the following:

Theorem A. *Let Γ be a convex co-compact group of a $\text{CAT}(-1)$ space X and let $\rho : \Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ be an irreducible convex representation with $d \geq 3$. Then*

$$\alpha h_{\rho} \leq h_{\Gamma} \text{ and } \alpha H_{\rho} \leq h_{\Gamma},$$

when $\xi : \text{L}_{\Gamma} \rightarrow \mathbb{P}(\mathbb{R}^d)$ is α -Hölder.

Remark that the dimension d of \mathbb{R}^d does not appear in the inequality.

Consider $\text{Ad} : \text{PGL}(d, \mathbb{R}) \rightarrow \text{PGL}(\mathfrak{sl}(d, \mathbb{R}))$ the Adjoint representation. If $\rho : \Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ is an irreducible convex representation then $\text{Ad } \rho : \Gamma \rightarrow \text{PGL}(\mathfrak{sl}(d, \mathbb{R}))$ is not necessarily irreducible but there is a natural subspace $V_{\rho} \subset \mathfrak{sl}(d, \mathbb{R})$ such that

$$A_{\rho} = \text{Ad } \rho|_{V_{\rho}} : \Gamma \rightarrow \text{PGL}(V_{\rho})$$

is again irreducible and convex (see Lemma 4.7). In the sequel, we will refer to A_{ρ} as the *irreducible adjoint representation* of ρ .

A simple computation shows that the Hilbert entropy of ρ is related to the spectral entropy of A_{ρ} , namely $H_{\rho} = 2h_{A_{\rho}}$. Nevertheless, applying this relation to the first inequality in Theorem A gives the bad upper bound $\alpha H_{\rho} \leq 2h_{\Gamma}$.

Examples. There are three examples of irreducible convex representations of Γ of particular interest.

Recall that the linear subgroup $\text{PSO}(1, k)$ of projective transformations preserving a bilinear form of signature $(1, k)$ is isomorphic to the isometry group $\text{Isom } \mathbb{H}^k$ of the hyperbolic space. Throughout this work we shall refer to the representation $\bar{\phi} : \text{Isom } \mathbb{H}^k \rightarrow \text{PSO}(1, k)$ (or any of its conjugates $g\bar{\phi}g^{-1}$ with $g \in \text{PGL}(k+1, \mathbb{R})$) as the *Klein model* of \mathbb{H}^k .

Remark 1.3. The Klein model of \mathbb{H}^k induces an equivariant map $\partial_{\infty}\mathbb{H}^k \rightarrow \mathbb{P}(\mathbb{R}^{k+1})$. This equivariant map is a bi-Lipschitz homeomorphism onto its image.

Benoist Representations: If $\rho : \Gamma \rightarrow \text{PGL}(k+1, \mathbb{R})$ preserves a proper open convex set Ω_{ρ} of $\mathbb{P}(\mathbb{R}^{k+1})$ and $\rho\Gamma \backslash \Omega_{\rho}$ is compact, then ρ is said to be a *Benoist representation*. Results from Benoist [4] imply that Benoist representations are irreducible convex representations (see [20] for details).

The Hilbert entropy of ρ is the topological entropy of the geodesic flow of $\rho\Gamma \backslash \Omega_{\rho}$ associated to the Hilbert metric. Crampon [12] proved that the Hilbert entropy H_{ρ} verifies $H_{\rho} \leq k-1 = \dim \partial\Omega_{\rho}$, and equality holds only when Ω_{ρ} is an ellipsoid, i.e. Γ acts co-compactly on \mathbb{H}^k and ρ extends to the Klein model of \mathbb{H}^k .

Convex co-compact groups in \mathbb{H}^k : Consider a convex co-compact group $\phi : \Gamma \rightarrow \text{Isom } \mathbb{H}^k$. The composition of ϕ with the Klein model of \mathbb{H}^k gives rise to a convex representation $\phi' : \Gamma \rightarrow \text{PGL}(k+1, \mathbb{R})$.

Remark that in this setting, $\phi\Gamma$ is Zariski dense in $\text{Isom } \mathbb{H}^k$ if and only if the action of $\phi\Gamma$ on \mathbb{H}^k does not have an invariant totally geodesic copy of \mathbb{H}^{k-1} . If this is the case, the convex representation $\phi'\Gamma$ is irreducible.

An easy computation shows that the spectral entropy of ϕ' , and the Hilbert entropy, coincide with the topological entropy of the geodesic flow of $\phi\Gamma\backslash\mathbb{H}^k$, which in turn coincides with the Hausdorff dimension of the limit set $L_{\phi\Gamma}$ on $\partial_\infty\mathbb{H}^k$, (Sullivan [22]).

Assume now that $\Gamma = \pi_1\mathbf{M}$ is the fundamental group of a closed k -dimensional hyperbolic manifold, it is well known that $h_\Gamma = k - 1$. Consider now a convex co-compact action $\phi : \pi_1\mathbf{M} \rightarrow \text{Isom } \mathbb{H}^n$ with $n \geq k$ as we explained before, Bourdon states that $h_\phi \geq k - 1$.

In light of the last examples one sees that a deformation of

$$\pi_1\mathbf{M} \rightarrow \text{Isom } \mathbb{H}^k \rightarrow \text{PGL}(k+1, \mathbb{R})$$

decreases Hilbert's entropy, but on the contrary, a deformation of

$$\pi_1\mathbf{M} \rightarrow \text{Isom } \mathbb{H}^k \rightarrow \text{Isom } \mathbb{H}^n$$

increases Hilbert's entropy. As a conclusion the Hilbert entropy of a convex representation of $\pi_1\mathbf{M}$ may be greater or smaller than $\dim \mathbf{M} - 1$, nevertheless the quantity $\alpha\mathbf{H}$ has to remain bounded by this number. Theorem A is thus optimal in this generality.

Hitchin representations and small deformations of exterior products:

Consider Σ a closed oriented hyperbolic surface and say that a representation $\rho : \pi_1\Sigma \rightarrow \text{PSL}(d, \mathbb{R})$ is *Fuchsian* if it factors as

$$\rho = \tau_d \circ \mathbf{f}$$

where $\tau_d : \text{PSL}(2, \mathbb{R}) \rightarrow \text{PSL}(d, \mathbb{R})$ the irreducible linear action (unique modulo conjugation) of $\text{PSL}(2, \mathbb{R})$ on \mathbb{R}^d and $\mathbf{f} : \pi_1\Sigma \rightarrow \text{PSL}(2, \mathbb{R})$ is a choice of a hyperbolic metric on Σ . A *Hitchin component* of $\text{PSL}(d, \mathbb{R})$ is a connected component of

$$\text{hom}(\pi_1\Sigma, \text{PSL}(d, \mathbb{R})) = \{\text{morphisms } \rho : \pi_1\Sigma \rightarrow \text{PSL}(d, \mathbb{R})\}$$

containing a Fuchsian representation. As Hitchin [15] proves, representations in the Hitchin component are irreducible.

Recall that a (complete) *flag* of \mathbb{R}^d is a collection of subspaces $\{V_i\}_{i=0}^d$ such that $V_i \subset V_{i+1}$ and $\dim V_i = i$. The space of flags is denoted \mathcal{F} . Two flags $\{V_i\}$ and $\{W_i\}$ are in *general position* if for every i one has

$$V_i \oplus W_{d-i} = \mathbb{R}^d.$$

Labourie [17] proves that if $\rho : \pi_1\Sigma \rightarrow \text{PSL}(d, \mathbb{R})$ is a representation in a Hitchin component then there exists a ρ -equivariant Hölder-continuous map $\zeta : \partial_\infty\pi_1\Sigma \rightarrow \mathcal{F}$. Moreover the flags $\zeta(x)$ and $\zeta(y)$ are in general position when $x, y \in \partial_\infty\pi_1\Sigma$ are distinct.

Considering thus $\xi := \zeta_1$ the first coordinate of ζ , and $\xi^* := \zeta_d$ the last coordinate of ζ , one obtains an irreducible convex representation. Moreover, let $\Lambda^n\mathbb{R}^d$ be the n -th exterior power of \mathbb{R}^d . An n -dimensional subspace is sent to a line on $\Lambda^n\mathbb{R}^d$, hence Labourie's theorem implies that the composition $\Lambda^n\rho : \pi_1\Sigma \rightarrow \text{PSL}(\Lambda^n\mathbb{R}^d)$ is again convex.

Finally, if ρ is Zariski dense on $\text{PGL}(d, \mathbb{R})$, then $\Lambda^n\rho$ is irreducible. Guichard and Wienhard [14] have shown that convex irreducible representations of hyperbolic groups form an open set on the set of representations. Hence small deformations of $\Lambda^n\rho$ are still irreducible and convex.

Remark 1.4. Labourie's statement implies that if $\rho : \pi_1 \Sigma \rightarrow \mathrm{PGL}(d, \mathbb{R})$ is a Hitchin representation, then the image $\xi(\partial_\infty \pi_1 \Sigma)$ is a curve of class C^1 (even though the map ξ is only Hölder). Hence, either entropy of ρ cannot be interpreted as the Hausdorff dimension of $\xi(\partial_\infty \pi_1 \Sigma)$. For example, if ρ is Fuchsian, then an easy computation shows that $h_\rho = H_\rho = 2/(d-1)$, even though the limit curve is a polynomial.

1.1. Equality. For a convex representation $\rho : \Gamma \rightarrow \mathrm{PGL}(d, \mathbb{R})$ denote by

$$\alpha_\rho = \sup\{\alpha \in (0, 1] : \xi : L_\Gamma \rightarrow \mathbb{P}(\mathbb{R}^d) \text{ is } \alpha\text{-Hölder}\}$$

the “best” Hölder exponent of the equivariant map ξ for a visual metric on L_Γ . Remark that ξ is not necessarily α_ρ -Hölder.

A key tool to study consequences of equality in Theorem A is the *cross ratio* associated to a convex representation introduced by Labourie [18]. Given a convex representation ρ define $\mathbf{b}_\rho : (\partial_\infty \Gamma)^{(4)} \rightarrow \mathbb{R}$ by

$$\mathbf{b}_\rho(x, y, z, t) = \frac{\varphi(u)}{\psi(u)} \frac{\psi(v)}{\varphi(v)} \quad (2)$$

where $\varphi \in \xi^*(x)$, $\psi \in \xi^*(z)$, $u \in \xi(y)$ and $v \in \xi(t)$. Remark that the result does not depend on the choice of φ , ψ , u and v made.

Theorem B (Spectral entropy rigidity). *Let Γ be a convex-co-compact group of \mathbb{H}^k that does not preserve a totally geodesic copy of \mathbb{H}^{k-1} , and consider a convex irreducible representation $\rho : \Gamma \rightarrow \mathrm{PGL}(d, \mathbb{R})$ with $d \geq 3$ such that equality*

$$\alpha_\rho h_\rho = h_\Gamma$$

holds. Assume also that $\mathbf{b}_\rho \geq 0$. Then $d = k + 1$, $\alpha_\rho = 1$ and ρ extends to $\bar{\rho} : \mathrm{Isom} \mathbb{H}^k \rightarrow \mathrm{PGL}(k + 1, \mathbb{R})$ as the Klein model of \mathbb{H}^k .

Remark that if $k \geq 3$ then the condition $\mathbf{b}_\rho \geq 0$ is a necessary condition for the Theorem to hold.

A weaker statement holds for Hilbert's entropy. Recall we have defined the adjoint irreducible representation of $\mathbf{A}_\rho : \Gamma \rightarrow \mathrm{PGL}(V_\rho)$ of a given ρ , as the restriction $\mathrm{Ad} \rho|_{V_\rho}$ where $V_\rho \subset \mathfrak{sl}(d, \mathbb{R})$. A slight modification of the proof of Theorem B gives the following.

Corollary 1.5 (Hilbert entropy rigidity). *Let Γ be a convex-co-compact group of \mathbb{H}^k that does not preserve a totally geodesic copy of \mathbb{H}^{k-1} , and consider a convex irreducible representation $\rho : \Gamma \rightarrow \mathrm{PGL}(d, \mathbb{R})$ with $d \geq 3$ such that equality*

$$\alpha_\rho H_\rho = h_\Gamma$$

holds. Assume also that $\mathbf{b}_\rho \geq 0$. Then $V_\rho = \mathfrak{so}(1, k)$ and the adjoint irreducible representation $\mathbf{A}_\rho : \Gamma \rightarrow \mathrm{PGL}(\mathfrak{so}(1, k))$ extends to $\overline{\mathbf{A}_\rho} : \mathrm{Isom} \mathbb{H}^k \rightarrow \mathrm{PGL}(\mathfrak{so}(1, k))$ as the adjoint representation of the Klein model of \mathbb{H}^k .

If Γ is a co-compact group in \mathbb{H}^2 then the fact that the cross ratio \mathbf{b}_ρ is non negative is consequence of the equality $\alpha_\rho h_\rho = h_\Gamma$:

Corollary 1.6. *Consider Σ a closed orientable hyperbolic surface and an irreducible convex representation $\rho : \pi_1 \Sigma \rightarrow \mathrm{PGL}(d, \mathbb{R})$ with $d \geq 3$ such that $\alpha_\rho h_\rho = 1$, then $d = 3$, $\alpha_\rho = 1$ and ρ extends to the Klein model of \mathbb{H}^2 .*

The proofs of Theorem B and Corollaries 1.5 and 1.6 are very similar and postponed to Section 11.

There are other several situations where the condition $\mathbf{b}_\rho \geq 0$ is automatically satisfied:

Corollary 1.7. *Let Γ be a convex-co-compact subgroup of \mathbb{H}^k $k \geq 2$ that does not preserve a totally geodesic copy of \mathbb{H}^{k-1} . Denote h_Γ the Hausdorff dimension of L_Γ . Consider a convex co-compact action $\rho : \Gamma \rightarrow \text{Isom } \mathbb{H}^n$ $n \geq 3$. If $\alpha_\rho h_\rho = h_\Gamma$ then ξ extends to $\bar{\xi} : \mathbb{H}^k \rightarrow \mathbb{H}^n$ as a ρ -equivariant isometric embedding.*

This is to say, $\rho\Gamma$ preserves a totally geodesic copy of \mathbb{H}^k in \mathbb{H}^n and moreover, the action of $\rho\Gamma$ on this geodesic copy is the departing action of Γ on \mathbb{H}^k .

Proof. Follows directly from Theorem B and the fact that the cross ratio $\mathbb{B}_{\mathbb{H}^n}$ of \mathbb{H}^n is non negative, see Section 6. \square

Definition 1.8. Assume now that $\partial_\infty \Gamma$ is connected and that it is not the circle, we will then say that Γ *splits over a virtual \mathbb{Z}* if Γ verifies either

- $\Gamma = A *_C B$ where A and B are non trivial groups and C is virtually \mathbb{Z} or,
- Γ is an HNN extension over a virtual \mathbb{Z} .

For example, the fundamental group of a closed hyperbolic manifold of dimension ≥ 3 or a Kleinian group of \mathbb{H}^3 having as boundary a Sierpiński carpet, do not split over a virtual \mathbb{Z} (Bowditch [8]).

Lemma 1.9 (See Lemma 6.3). *Let Γ be a convex-co-compact group of \mathbb{H}^k that does not preserve a totally geodesic copy of \mathbb{H}^{k-1} , such that $\partial_\infty \Gamma$ is connected and not homeomorphic to the circle. Assume that Γ does not split over a virtual \mathbb{Z} and consider a convex irreducible representation $\rho : \Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ then $\mathbf{b}_\rho \geq 0$.*

Hence, for such groups Theorem B and Corollary 1.5 hold without the assumption $\mathbf{b}_\rho \geq 0$.

1.2. Statements for hyperconvex representations. The fact that equality in Corollary 1.6 can only hold for a representation $\rho : \pi_1 \Sigma \rightarrow \text{PSL}(3, \mathbb{R})$ suggests that the upper bound for $\alpha_\rho h_\rho$ is not optimal for Hitchin representations on $\text{PSL}(d, \mathbb{R})$, say. We will now focus on improving the bound when more information on the representation ρ is given. Let G be a real algebraic semi-simple Lie group, P be a minimal parabolic subgroup of G , and write $\mathcal{F} = G/P$ for the *Furstenberg boundary* of the symmetric space of G .

Let K be a maximal compact subgroup of G , let τ be the Cartan involution on \mathfrak{g} for which the set $\text{fix } \tau$ is K 's Lie algebra, consider $\mathfrak{p} = \{v \in \mathfrak{g} : \tau v = -v\}$ and \mathfrak{a} a maximal abelian subspace contained in \mathfrak{p} . Let Σ be the set of (restricted) roots of \mathfrak{a} on \mathfrak{g} . Fix \mathfrak{a}^+ a closed Weyl chamber and let Σ^+ a system of positive roots on Σ associated to \mathfrak{a}^+ , denote by Π the set of simple roots associated to the choice Σ^+ .

The space \mathcal{F} can be embedded in a product of projective spaces $\prod_{\theta \in \Pi} \mathbb{P}(V_\theta)$ (see Section 9), we will consider the metric on \mathcal{F} induced by this embedding.

The product $\mathcal{F} \times \mathcal{F}$ has a unique open G -orbit denoted by $\mathcal{F}^{(2)}$. For example, if $G = \text{PGL}(d, \mathbb{R})$ then \mathcal{F} is the space of complete flags of \mathbb{R}^d , and $\mathcal{F}^{(2)}$ is the space of flags in general position.

Definition 1.10. We say that a representation $\rho : \Gamma \rightarrow G$ is *hyperconvex* if there exists a Hölder-continuous equivariant mapping $\zeta : \partial_\infty \Gamma \rightarrow \mathcal{F}$ such that the pair $(\zeta(x), \zeta(y))$ belongs to $\mathcal{F}^{(2)}$ whenever $x, y \in \partial_\infty \Gamma$ are distinct.

Hyperconvex representations on $\mathrm{PGL}(d, \mathbb{R})$ are of course convex. As we explained before, Labourie [17] has shown that representations in a Hitchin component are hyperconvex.

The *barycenter* of the Weyl chamber \mathfrak{a}^+ is the half line contained in \mathfrak{a}^+ determined by

$$\mathrm{bar}_{\mathfrak{a}^+} = \{a \in \mathfrak{a}^+ : \theta_1(a) = \theta_2(a) \text{ for every pair } \theta_1, \theta_2 \in \Pi\}.$$

We will say that $g \in G$ is \mathbb{R} -*regular* if it is diagonalizable over \mathbb{R} , *elliptic* if it is contained in a compact subgroup of G , or *unipotent* if all its eigenvalues are equal to 1.

Recall that Jordan's decomposition states that every $g \in G$ can be written as a product $g = g_e g_h g_u$ where $g_e, g_h, g_u \in G$ commute, g_e is elliptic, g_h is \mathbb{R} -regular and g_u is unipotent.

For $g \in G$ denote by $\lambda(g) \in \mathfrak{a}^+$ its *Jordan projection*, this is the unique element on \mathfrak{a}^+ such that $\exp \lambda(g)$ is conjugated to the \mathbb{R} -regular element on the Jordan decomposition of g .

Again, Γ is a convex co-compact group of a $\mathrm{CAT}(-1)$ space X and h_Γ is the Hausdorff dimension of the limit set L_Γ on $\partial_\infty X$. For a hyperconvex representation $\rho : \Gamma \rightarrow G$ and a linear functional $\varphi \in \mathfrak{a}^*$ such that $\varphi|_{\mathfrak{a}^+} > 0$ we define the *entropy of ρ relative to φ* as

$$h_\varphi := \lim_{s \rightarrow \infty} \frac{\log \#\{[\gamma] \in [\Gamma] : \varphi(\lambda(\rho\gamma)) \leq s\}}{s}.$$

Proposition 1.11 ([20, Section 7]). *Let $\rho : \Gamma \rightarrow G$ a Zariski dense hyperconvex representation and consider $\varphi \in \mathfrak{a}^*$ such that $\varphi|_{\mathfrak{a}^+} - \{0\} > 0$, then $h_\varphi \in (0, \infty)$.*

For a hyperconvex representation we have a much better bound on the quantity αh_φ .

Theorem C. *Let $\rho : \Gamma \rightarrow G$ be a Zariski dense hyperconvex representation, and $\varphi \in \mathfrak{a}^*$ a linear functional such that $\varphi|_{\mathfrak{a}^+} - \{0\} > 0$ then*

$$\alpha h_\varphi \leq h_\Gamma \frac{\theta(\mathrm{bar}_{\mathfrak{a}^+})}{\varphi(\mathrm{bar}_{\mathfrak{a}^+})}$$

where $\theta \in \Pi$ is any simple root and α is the Hölder exponent of the equivariant map $\zeta : L_\Gamma \rightarrow \mathcal{F}$.

Remark that the direction of \mathfrak{a}^+ that gives the upper bound does not depend on the linear form φ .

In order to study rigidity when equality holds we need to restrict the functionals we consider. If χ is a restricted weight of G denote by $\Lambda_\chi : G \rightarrow \mathrm{PGL}(V_\chi)$ the irreducible proximal representation associated to it (Proposition 9.1). The composition $\Lambda_\chi \circ \rho : \Gamma \rightarrow \mathrm{PGL}(V_\chi)$ is a convex representation of Γ (Lemma 12.2), we will denote by b_χ the cross ratio defined on $\partial_\infty \Gamma$ associated to $\Lambda_\chi \circ \rho$ as in equation (2).

For a hyperconvex representation $\rho : \Gamma \rightarrow G$ denote by

$$\alpha_\rho = \sup\{\alpha \in (0, 1] : \zeta : L_\Gamma \rightarrow \mathcal{F} \text{ is } \alpha\text{-Hölder}\}.$$

Theorem D. *Let Γ be a convex-co-compact group of \mathbb{H}^k that does not preserve a totally geodesic copy of \mathbb{H}^{k-1} , and consider a Zariski dense hyperconvex representation $\rho : \Gamma \rightarrow G$. Consider a restricted weight χ of G and assume that*

$$\alpha_\rho h_\chi = h_\Gamma \frac{\theta(\text{bar}_{\mathfrak{a}^+})}{\chi(\text{bar}_{\mathfrak{a}^+})}$$

where $\theta \in \Pi$ is any simple root. Suppose also that $\mathbf{b}_\chi \geq 0$. Then $\text{Isom } \mathbb{H}^k$ is a factor of G and the following diagram commutes

$$\begin{array}{ccc} \Gamma & \xrightarrow{\rho} & G \\ & \searrow & \downarrow \\ & & \text{Isom } \mathbb{H}^k \end{array}$$

where the arrow $\Gamma \rightarrow \text{Isom } \mathbb{H}^k$ is the action of Γ on \mathbb{H}^k fixed at the beginning.

As for Theorem B, the condition $\mathbf{b}_\chi \geq 0$ is not needed when Γ does not split over a virtual \mathbb{Z} or when Γ act co-compactly on \mathbb{H}^2 . A slight modification of the proof of Theorem D gives the following.

Corollary 1.12. *Let Γ be a co-compact group of \mathbb{H}^2 and consider a Zariski dense hyperconvex representation $\rho : \pi_1 \Sigma \rightarrow G$. Suppose the equality*

$$\alpha_\rho h_\chi = \frac{\theta(\text{bar}_{\mathfrak{a}^+})}{\chi(\text{bar}_{\mathfrak{a}^+})}$$

holds for some restricted weight χ and any simple root θ . Then $\text{PSL}(2, \mathbb{R})$ is a factor of G .

Theorem C, Corollary 1.12 and a Theorem of Guichard (14.1 below) give the following corollary whose proof is postponed to the end of this article.

Corollary 1.13. *Let $\rho : \pi_1 \Sigma \rightarrow \text{PGL}(d, \mathbb{R})$ be a representation in the Hitchin component then*

$$\alpha_\rho h_\rho \leq \frac{2}{d-1} \text{ and } \alpha_\rho \mathbf{H}_\rho \leq \frac{2}{d-1}$$

and either equality holds only if $\rho = \tau_d \circ \mathbf{f}$, where $\tau_d : \text{PSL}(2, \mathbb{R}) \rightarrow \text{PSL}(d, \mathbb{R})$ is the irreducible representation and $\mathbf{f} : \pi_1 \Sigma \rightarrow \text{PSL}(2, \mathbb{R})$ is the departing action.

Remark that if $\zeta : \partial_\infty \pi_1 \Sigma \rightarrow \mathcal{F}$ is the equivariant map of a Hitchin representation then by definition, it is less (or equally) regular than the induced equivariant map $\xi = \zeta_1 : \partial_\infty \pi_1 \Sigma \rightarrow \mathbb{P}(\mathbb{R}^d)$. Hence, even though we obtain a much better bound on $\alpha_\rho h_\rho$ we do not know if this is produced by a decay of regularity of the map ζ .

1.3. The method. Let us explain the main idea for the proof of Theorem C for $G = \text{PGL}(d, \mathbb{R})$ (Theorem A is proved in a similar fashion).

A Cartan subspace of $G = \text{PGL}(d, \mathbb{R})$ is $\mathfrak{a} = \{(a_1, \dots, a_d) \in \mathbb{R}^d : a_1 + \dots + a_d = 0\}$, a Weyl chamber is $\mathfrak{a}^+ = \{(a_1, \dots, a_d) \in \mathfrak{a} : a_1 \geq \dots \geq a_d\}$, and the simple roots associated to \mathfrak{a}^+ are

$$\Pi = \{\theta_i(a_1, \dots, a_d) = a_i - a_{i+1} : i \in \{1, \dots, d-1\}\}.$$

The *walls* of the Weyl chamber \mathfrak{a}^+ are

$$W_i = \{a \in \mathfrak{a}^+ : a_i - a_{i+1} = 0\}.$$

Denote by $\lambda : \mathrm{PGL}(d, \mathbb{R}) \rightarrow \mathfrak{a}^+$ the Jordan projection, i.e.

$$\lambda(g) = (\lambda_1(g), \dots, \lambda_d(g))$$

are the logarithm of the modulus of the eigenvalues of g , counted with multiplicity and in decreasing order.

Consider, for example, the linear functional $\varphi : \mathfrak{a}^+ \rightarrow \mathbb{R}$ defined by

$$\varphi(a_1, \dots, a_d) = \frac{a_1 - a_d}{2}.$$

Remark that h_φ , the entropy relative to φ , is the Hilbert entropy H_ρ of ρ .

For an element $g \in \mathrm{PGL}(d, \mathbb{R})$ the number

$$\frac{\theta_i(\lambda(g))}{\varphi(\lambda(g))} = 2 \frac{\lambda_i(g) - \lambda_{i+1}(g)}{\lambda_1(g) - \lambda_d(g)}$$

measures, in some sense, how far the half line $\mathbb{R}_+ \lambda(g)$ is from the wall W_i .

The half line which is as far as possible from *all* the walls of \mathfrak{a}^+ is exactly the barycenter $\mathrm{bar}_{\mathfrak{a}^+}$, this line is determined by the equalities

$$a \in \mathfrak{a}^+ : a_1 - a_2 = a_2 - a_3 = \dots = a_{d-1} - a_d,$$

i.e. the barycenter is

$$\mathrm{bar}_{\mathfrak{a}^+} = \{((d-1)t, (d-3)t, \dots, (3-d)t, (1-d)t) : t \in \mathbb{R}_+\}.$$

For $a \in \mathrm{bar}_{\mathfrak{a}^+}$ one has

$$\frac{\theta_1(a)}{\varphi(a)} = 2 \frac{a_1 - a_2}{a_1 - a_d} = \frac{2}{d-1}.$$

Hence, the upper bound on Theorem C for $\mathrm{PGL}(d, \mathbb{R})$ is $2/(d-1)$.

Given a Zariski dense hyperconvex representation $\rho : \Gamma \rightarrow \mathrm{PGL}(d, \mathbb{R})$ we need to find an element $\gamma \in \Gamma$ such that

$$\alpha \frac{H_\rho}{h_\Gamma} \leq 2 \frac{\lambda_i(\rho\gamma) - \lambda_{i+1}(\rho\gamma)}{\lambda_1(\rho\gamma) - \lambda_d(\rho\gamma)}$$

for all $i \in \{1, \dots, d-1\}$. This is done using the Thermodynamic Formalism for convex representations developed on [20], together with some Linear Algebra of Benoist [4].

Let us explain now the main idea of Theorem B.

Consider an irreducible convex representation $\rho : \Gamma \rightarrow \mathrm{PGL}(d, \mathbb{R})$ where Γ is convex co-compact on \mathbb{H}^k and does not preserve a totally geodesic copy of \mathbb{H}^{k-1} (as explained before, this is equivalent to Γ being Zariski dense in $\mathrm{Isom} \mathbb{H}^k$). Remark we are not making any assumption on the Zariski closure of $\rho\Gamma$.

If equality $\alpha h_\rho = h_\Gamma$ holds, the Thermodynamic Formalism will imply the following relation for every non torsion $\gamma \in \Gamma$:

$$\lambda_1(\rho\gamma) = c|\gamma|$$

for some $c \in \mathbb{R}_+^*$ and where $|\gamma|$ is the translation length of $\gamma \in \Gamma$.

The modulus of the cross ratio \mathbf{b}_ρ is uniquely determined by the spectral radii $\{\lambda_1(\rho\gamma) : \gamma \in \Gamma\}$ (see Lemma 7.3), hence the cross ratio $|\mathbf{b}_\rho|$ of the representation ρ and the cross ratio $\mathbb{B}_{\mathbb{H}^k}^c|_{L_\Gamma}$ coincide, where $\mathbb{B}_{\mathbb{H}^k}|_{L_\Gamma}$ is the natural cross ratio on $\partial_\infty \mathbb{H}^k$ restricted to the limit set L_Γ of Γ on $\partial_\infty \mathbb{H}^k$.

Since \mathbf{b}_ρ is positive we have $\mathbf{b}_\rho = \mathbb{B}_{\mathbb{H}^k}^c|_{L_\Gamma}$. In order to extend $\rho : \Gamma \rightarrow \mathrm{PGL}(d, \mathbb{R})$ to $\bar{\rho} : \mathrm{Isom} \mathbb{H}^k \rightarrow \mathrm{PGL}(d, \mathbb{R})$ we will use a Theorem of Labourie 6.6 which asserts that, when a cross ratio has finite rank it is induced by a convex representation.

We need then to show that the cross ratio $\mathbb{B}_{\mathbb{H}^k}^c$, defined on all $\partial_\infty \mathbb{H}^k$, has finite rank. In order to do so we will use some techniques from Benoist [2]. The fact that $\text{Isom } \mathbb{H}^k$ is a real algebraic group is crucial in this step.

Acknowledgments. The author is extremely thankful to Martin Bridgeman, Dick Canary, Francois Labourie, Alejandro Passeggi, Rafael Potrie and Jean-François Quint for useful discussions. He would like to particularly thank Yves Benoist and Matias Carrasco for discussions that considerably improved the statements of this work, and Qiongleng Li for pointing out an error on the first version of this paper.

2. REPARAMETRIZATIONS AND THERMODYNAMIC FORMALISM

Let X be a compact metric space and let $\phi = \{\phi_t : X \rightarrow X\}_{t \in \mathbb{R}}$ be a continuous flow on X without fixed points. Consider a positive continuous function $f : X \rightarrow \mathbb{R}_+^*$ and define $\kappa : X \times \mathbb{R} \rightarrow \mathbb{R}$ by

$$\kappa(x, t) = \int_0^t f(\phi_s(x)) ds. \quad (3)$$

The function κ has the cocycle property $\kappa(x, t + s) = \kappa(\phi_t x, s) + \kappa(x, t)$ for every $t, s \in \mathbb{R}$ and $x \in X$.

Since $f > 0$ and X is compact f has a positive minimum and $\kappa(x, \cdot)$ is an increasing homeomorphism of \mathbb{R} . We then have an inverse $\alpha : X \times \mathbb{R} \rightarrow \mathbb{R}$ such that

$$\alpha(x, \kappa(x, t)) = \kappa(x, \alpha(x, t)) = t \quad (4)$$

for every $(x, t) \in X \times \mathbb{R}$.

Definition 2.1. The *reparametrization* of ϕ by f is the flow $\psi = \psi^f = \{\psi_t : X \rightarrow X\}_{t \in \mathbb{R}}$ defined by $\psi_t(x) = \phi_{\alpha(x, t)}(x)$. If f is Hölder-continuous we will say that ψ is a Hölder reparametrization of ϕ .

We say that a function $U : X \rightarrow \mathbb{R}$ is C^1 *in the direction of the flow* ϕ if for every $p \in X$ the function $t \mapsto U(\phi_t(p))$ is of class C^1 and the function

$$p \mapsto \left. \frac{\partial}{\partial t} \right|_{t=0} U(\phi_t(p))$$

is continuous. Two Hölder-continuous functions $f, g : X \rightarrow \mathbb{R}$ are then said to be *Livšic-cohomologous* if there exists a continuous $U : X \rightarrow \mathbb{R}$, of class C^1 in the direction of the flow, such that for all $p \in X$ one has

$$f(p) - g(p) = \left. \frac{\partial}{\partial t} \right|_{t=0} U(\phi_t(p)).$$

Remark 2.2. When two continuous functions $f, g : X \rightarrow \mathbb{R}_+^*$ are Livšic-cohomologous the reparametrization of ϕ by f is conjugated to the reparametrization by g , i.e. there exists a homeomorphism $h : X \rightarrow X$ such that for all $p \in X$ and $t \in \mathbb{R}$

$$h(\psi_t^f p) = \psi_t^g(hp).$$

Let ψ be the reparametrization of ϕ by $f : X \rightarrow \mathbb{R}_+^*$. If τ is a periodic orbit of ϕ then the period of τ for ψ is

$$\int_\tau f = \int_0^{p(\tau)} f(\phi_s(x)) ds \quad (5)$$

where $p(\tau)$ is the period of τ for ϕ and $x \in \tau$.

If m is a ϕ -invariant probability measure on X then the probability measure $m^\#$ defined by

$$\frac{dm^\#}{dm}(\cdot) = f(\cdot) / \int f dm$$

is ψ -invariant. This relation between invariant probability measures induces a bijection and Abramov [1] relates the corresponding metric entropies:

$$h(\psi, m^\#) = h(\phi, m) / \int f dm. \quad (6)$$

Denote by \mathcal{M}^ϕ the set of ϕ -invariant probability measures. The *pressure* of a continuous function $f : X \rightarrow \mathbb{R}$ is defined by

$$P(\phi, f) = \sup_{m \in \mathcal{M}^\phi} h(\phi, m) + \int_X f dm.$$

A probability m such that the supremum is attained is called an *equilibrium state* of f . An equilibrium state for $f \equiv 0$ is called a *probability of maximal entropy*, its entropy is called the *topological entropy* of ϕ and denoted $h_{\text{top}}(\phi)$.

Lemma 2.3 ([20, Section 2]). *Let ψ be the reparametrization of ϕ by $f : X \rightarrow \mathbb{R}_+^*$, and assume that $h_{\text{top}}(\psi)$ is finite. Then $m \mapsto m^\#$ induces a bijection between the set of equilibrium states of $-h_{\text{top}}(\psi)f$ and the set of probability measures of maximal entropy of ψ .*

2.1. Topological Anosov flows. For $\varepsilon > 0$ one defines the *local stable set* of x by

$$W_\varepsilon^s(x) = \{y \in X : d(\phi_t x, \phi_t y) \leq \varepsilon \forall t > 0 \text{ and } d(\phi_t x, \phi_t y) \rightarrow 0 \text{ as } t \rightarrow \infty\}$$

and the local unstable set by

$$W_\varepsilon^u(x) = \{y \in X : d(\phi_{-t} x, \phi_{-t} y) \leq \varepsilon \forall t > 0 \text{ and } d(\phi_{-t} x, \phi_{-t} y) \rightarrow 0 \text{ as } t \rightarrow \infty\}.$$

Definition 2.4. We will say that ϕ is a *topological Anosov flow* if the following holds:

- There exist positive constants C , λ and ε such that for every $x \in X$, every $y \in W_\varepsilon^s(x)$ and every $t > 0$ one has

$$d(\phi_t(x), \phi_t(y)) \leq C e^{-\lambda t}$$

and such that for every $y \in W_\varepsilon^u(x)$ one has

$$d(\phi_{-t}(x), \phi_{-t}(y)) \leq C e^{-\lambda t}.$$

- For every x there exists a neighborhood U of x such that if $y, z \in U$ then $W_\varepsilon^s(z) \cap W_\varepsilon^u(y)$ consists of at most one point and such that, for ε and $\delta > 0$ small enough, the map $W_\varepsilon^s(x) \times W_\varepsilon^u(x) \times [-\delta, \delta] \rightarrow X$ defined by

$$(y, z, t) \mapsto \phi_t(W_\varepsilon^u(y) \cap W_\varepsilon^s(z))$$

is a well defined homeomorphism onto some neighborhood of x .

A flow is said to be *transitive* if it has a dense orbit. Anosov's closing Lemma is a standard dynamical tool in hyperbolic dynamics.

Theorem 2.5 (Anosov's closing Lemma c.f. Shub [21]). *Let ϕ be transitive topological Anosov flow, then periodic orbits are dense in the set of ergodic invariant probability measures of ϕ .*

The following is standard in the study of Ergodic Theory of Anosov flows.

Proposition 2.6 (Bowen-Ruelle [9]). *Let ϕ be a transitive topological Anosov flow. Then given a Hölder-continuous function $f : X \rightarrow \mathbb{R}$ there exists a unique equilibrium state for f , moreover, the equilibrium state is ergodic. If two functions have the same equilibrium state then their difference is Livšic-cohomologous to a constant.*

We will need the following immediate Lemma.

Lemma 2.7. *Let ϕ be a topological Anosov flow on X and $f : X \rightarrow \mathbb{R}_+$ a positive Hölder-continuous function. Denote by*

$$h_f = \lim_{t \rightarrow \infty} \frac{1}{t} \log \# \{ \tau\text{-periodic} : \int_{\tau} f \leq t \},$$

then

$$\frac{h(\phi, m_{-h_f f})}{h_f} = \int f dm_{-h_f f}.$$

Proof. Let ϕ^f be the reparametrization of ϕ by f . The flow ϕ^f is still a topological Anosov flow and hence its topological entropy can then be computed as the exponential growth rate of its periodic orbits, i.e. the topological entropy of ϕ^f is h_f (recall equation (5)). The Lemma finishes applying Lemma 2.3 and Abramov's formula (6). \square

3. CAT(−1) SPACES

The standard reference for this section is Bourdon [6]. Consider a CAT(−1) space X and $\partial_\infty X$ its visual boundary. The *Busseman function* of X , $B : \partial_\infty X \times X \times X \rightarrow \mathbb{R}$ is defined by

$$B(z, p, q) = B_z(p, q) = \lim_{s \rightarrow \infty} d_X(p, \sigma(s)) - d_X(q, \sigma(s))$$

where $\sigma : [0, \infty) \rightarrow X$ is any geodesic ray such that $\sigma(\infty) = z$.

Denote by

$$\partial_\infty^{(2)} X = \partial_\infty X \times \partial_\infty X - \{(x, x) : x \in \partial_\infty X\}$$

and fix a point $o \in X$. The *Gromov product* of X based on o , $[\cdot, \cdot]_o : \partial_\infty^{(2)} X \rightarrow \mathbb{R}$, is defined by

$$[x, y]_o = \frac{1}{2}(B_x(o, p) + B_y(o, p))$$

where p is any point in the geodesic joining x and y . Remark that $[x, y]_o \rightarrow \infty$ as y approaches x . The *visual metric* based on o on $\partial_\infty X$ is defined by $\delta_o(x, y) = e^{-[x, y]_o}$. Since X is CAT(−1) this is in fact a distance on $\partial_\infty X$.

For $\gamma \in \text{Isom } X$ denote by $|\gamma|$ its translation length

$$|\gamma| = \inf_{p \in X} d_X(p, \gamma p).$$

If γ is hyperbolic then one has $|\gamma| = B_{\gamma_+}(\gamma^{-1}o, o)$ for any $o \in X$ where γ_+ is the attractor of γ on $\partial_\infty X$.

The following Lemma shows that the translation length of a hyperbolic element can be directly computed from its action on the visual boundary of X .

Lemma 3.1. *Consider a hyperbolic element $\gamma \in \text{Isom } X$ then for any $x \in \partial_\infty X - \{\gamma_-\}$ one has*

$$\lim_{n \rightarrow \infty} \frac{\log \delta_o(\gamma^n x, \gamma_+)}{n} = -|\gamma|.$$

Proof. This is standard (Yue [24]). Fix two points $x, z \in \partial_\infty X$, then for every $\gamma \in \text{Isom } X$ one has

$$\delta_o(\gamma z, \gamma x) = e^{\frac{1}{2}(B_{\gamma z}(\gamma o, o) + B_{\gamma x}(\gamma o, o))} \delta_o(z, x).$$

Hence, for each δ there exists a neighborhood V of z such that for every $x \in V$ one has

$$1 - \delta \leq \frac{\delta_o(\gamma z, \gamma x)}{\delta_o(z, x)} e^{-B_{\gamma z}(\gamma o, o)} \leq 1 + \delta.$$

Assume now that γ is a hyperbolic element and that $z = \gamma_+$. Fix δ and assume that $x \in V$, then one has

$$(1 - \delta)^n \leq \frac{\delta_o(\gamma_+, \gamma^n x)}{\delta_o(\gamma_+, x)} e^{-nB_{\gamma_+}(\gamma o, o)} \leq (1 + \delta)^n.$$

Taking logarithm and dividing by n one obtains the desired conclusion. If $x \notin V$ then a big enough power $\gamma^N x$ does (recall $x \neq \gamma_-$) and one repeats the argument. \square

For a discrete subgroup Γ of $\text{Isom } X$ denote by L_Γ its limit set on $\partial_\infty X$. Consider the space $\widetilde{\text{U}\Gamma}$ defined by

$$\{\sigma : (-\infty, \infty) \rightarrow X : \sigma \text{ is a complete geodesic with } \sigma(-\infty), \sigma(\infty) \in L_\Gamma\}.$$

The group Γ naturally acts on $\widetilde{\text{U}\Gamma}$ and we denote by $\text{U}\Gamma = \Gamma \backslash \widetilde{\text{U}\Gamma}$ its quotient. We will say that Γ is *convex co-compact* if the space $\text{U}\Gamma$ is compact.

Remark 3.2. Throughout this work we will fix a convex co-compact action of Γ on X , hence we allow ourselves to naturally identify L_Γ to $\partial_\infty \Gamma$ and to refer to the space $\text{U}\Gamma$ as only depending on Γ .

The space $\text{U}\Gamma$ is naturally equipped with a flow $\phi = \{\phi_t : \text{U}\Gamma \rightarrow \text{U}\Gamma\}_{t \in \mathbb{R}}$ simply by changing the parametrization of a given complete geodesic. This is called *the geodesic flow of Γ*

The following Theorem relates this section to the preceding one:

Theorem 3.3 (c.f. Bourdon [6]). *Let Γ be a convex-co-compact group of X . Then the geodesic flow of Γ is a topological Anosov flow. The topological entropy of the geodesic flow is hence*

$$h_\Gamma = \lim_{s \rightarrow \infty} \frac{\log \#\{[\gamma] \in [\Gamma] \text{ non torsion} : |\gamma| \leq s\}}{s}.$$

3.1. Hölder cocycles. We shall now study Hölder cocycles on $\partial_\infty \Gamma$. The main references for this subsection are Ledrappier [19] and [20, Section 5].

Definition 3.4. A *Hölder cocycle* is a function $c : \Gamma \times \partial_\infty \Gamma \rightarrow \mathbb{R}$ such that

$$c(\gamma_0 \gamma_1, x) = c(\gamma_0, \gamma_1 x) + c(\gamma_1, x)$$

for any $\gamma_0, \gamma_1 \in \Gamma$ and $x \in \partial_\infty \Gamma$, and where $c(\gamma, \cdot)$ is a Hölder map for every $\gamma \in \Gamma$ (the same exponent is assumed for every $\gamma \in \Gamma$).

Given a Hölder cocycle c and a non torsion element $\gamma \in \Gamma$ we define the *period* of γ for c by

$$\ell_c(\gamma) = c(\gamma, \gamma_+)$$

where γ_+ is the attractive fixed point of γ on $\partial_\infty \Gamma$. The cocycle property implies that the period of an element γ only depends on its conjugacy class $[\gamma] \in [\Gamma]$.

We shall be interested in cocycles whose periods are positive, i.e. such that $\ell_c(\gamma) > 0$ for every non torsion $\gamma \in \Gamma$. The *entropy*[†] of such cocycle is defined by:

$$h_c = \limsup_{t \rightarrow \infty} \frac{1}{t} \log \# \{ [\gamma] : \ell_c(\gamma) \leq t \} \in \mathbb{R}_+ \cup \{ \infty \}.$$

Theorem 3.5 ([20, Section 3]). *Let c be a Hölder cocycle such that $h_c \in (0, \infty)$, then there exists a positive Hölder-continuous function $f : \text{U}\Gamma \rightarrow \mathbb{R}_+^*$ such that*

$$\int_{[\gamma]} f = \ell_c(\gamma)$$

for every non torsion $[\gamma]$.

4. CONVEX REPRESENTATIONS

Let Γ be a convex co-compact group of a $\text{CAT}(-1)$ space.

Definition 4.1. A representation $\rho : \Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ is *convex* if there exists a ρ -equivariant Hölder-continuous map

$$(\xi, \xi^*) : \partial_\infty \Gamma \rightarrow \mathbb{P}(\mathbb{R}^d) \times \mathbb{P}(\mathbb{R}^{d*}),$$

such that $\mathbb{R}^d = \xi(x) \oplus \ker \xi^*(y)$ whenever $x \neq y$.

Lemma 4.2. *Let $\rho : \Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ be a convex representation, then the action of $\rho\Gamma$ on $\langle \xi(\partial_\infty \Gamma) \rangle$ is irreducible.*

Proof. Consider $W \subset \langle \xi(\partial_\infty \Gamma) \rangle$ a $\rho\Gamma$ -invariant subspace. Consider $w \in W$ and write

$$w = \sum_{i=1}^k \alpha_i v_i$$

where $v_i \in \xi(x_i)$ for k -points $x_i \in \partial_\infty \Gamma$. Consider now some non torsion $\gamma \in \Gamma$ such that $\gamma_- \notin \{x_1, \dots, x_k\}$. We then have $\gamma^n x_i \rightarrow \gamma_+$ and hence $\mathbb{R}\rho\gamma^n(w) \rightarrow \xi(\gamma_+)$ in $\mathbb{P}(\mathbb{R}^d)$. Thus $\xi(\gamma_+) \in W$, since W is $\rho\Gamma$ -invariant one has

$$\xi(\partial_\infty \Gamma) = \xi(\overline{\Gamma \cdot \gamma_+}) \subset W.$$

This finishes the proof. □

We say that $g \in \text{PGL}(d, \mathbb{R})$ is *proximal* if it has a unique complex eigenvalue of maximal modulus, and its generalized eigenspace is one dimensional. This eigenvalue is necessarily real and its modulus is equal to $\exp \lambda_1(g)$. We will denote by g_+ the g -fixed line of \mathbb{R}^d consisting of eigenvectors of this eigenvalue, and g_- the g -invariant complement of g_+ (i.e. $\mathbb{R}^d = g_+ \oplus g_-$). The line g_+ is an attractor on $\mathbb{P}(\mathbb{R}^d)$ for the action of g and g_- is a repelling hyperplane.

Lemma 4.3 ([20, Section 3]). *Let $\rho : \Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ be a convex irreducible representation. Then for every non torsion element $\gamma \in \Gamma$, $\rho(\gamma)$ is proximal, $\xi(\gamma_+)$ is its attractive fixed line and $\xi^*(\gamma_-)$ is the repelling hyperplane. Consequently $\xi(x) \subset \xi^*(x)$ for every $x \in \partial_\infty \Gamma$.*

[†]In [20] this is called the exponential growth rate of the cocycle.

Fix now some norm $\|\cdot\|$ on \mathbb{R}^d . We define the Hölder cocycles $\beta_\rho, \bar{\beta}_\rho : \Gamma \times \partial_\infty \Gamma \rightarrow \mathbb{R}$ by

$$\beta_\rho(\gamma, x) = \log \frac{\|\rho(\gamma)v\|}{\|v\|} \text{ and } \bar{\beta}_\rho(\gamma, x) = \log \frac{\|\theta \circ \rho(\gamma^{-1})\|}{\|\theta\|}$$

for some non zero $v \in \xi(x)$, and a non zero functional $\theta \in \xi^*(x)$. Lemma 4.3 implies the following.

Lemma 4.4 ([20, Section 3]). *Assume ρ is convex and irreducible, then the period of a non torsion $\gamma \in \Gamma$ for β_ρ is $\beta_\rho(\gamma, \gamma_+) = \lambda_1(\rho\gamma)$, and the period $\bar{\beta}_\rho(\gamma, \gamma_+) = \lambda_1(\rho\gamma^{-1}) = -\lambda_d(\rho\gamma)$.*

Proposition 1.2 together with the last Lemma and Theorem 3.5 imply the following.

Corollary 4.5 ([20, Section 5]). *Let $\rho : \Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ be an irreducible convex representation, then there exists a positive Hölder-continuous function $f : \text{U}\Gamma \rightarrow \mathbb{R}_+^*$ such that*

$$\int_{[\gamma]} f = \lambda_1(\rho\gamma)$$

for every non torsion $[\gamma]$.

4.1. Adjoint representation. Given an irreducible convex representation $\rho : \Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ we will now show how the Adjoint representation $\text{Ad} : \text{PGL}(d, \mathbb{R}) \rightarrow \text{PGL}(\mathfrak{sl}(d, \mathbb{R}))$ induces again an irreducible convex representation \mathbf{A}_ρ such that

$$\lambda_1(\mathbf{A}_\rho\gamma) = \lambda_1(\rho\gamma) - \lambda_d(\rho\gamma).$$

This is standard.

Recall that the *adjoint representation* is defined by conjugation $\text{Ad}(g)(T) = gTg^{-1}$, where $T \in \mathfrak{sl}(d, \mathbb{R}) = \{\text{traceless endomorphisms of } \mathbb{R}^d\}$. Consider $\mathcal{F}_*(\mathbb{R}^d)$ the space of incomplete flags consisting of a line contained on a hyperplane,

$$\mathcal{F}_*(\mathbb{R}^d) = \{(v, \theta) \in \mathbb{P}(\mathbb{R}^d) \times \mathbb{P}(\mathbb{R}^{d*}) : \theta(v) = 0\}.$$

Given $(v, \theta) \in \mathcal{F}_*$ define $M(v, \theta) \in \mathbb{P}(\mathfrak{sl}(d, \mathbb{R}))$ by $M(v, \theta)(w) = \theta(w)v$ and define $\Phi(v, \theta) \in \mathbb{P}(\mathfrak{sl}(d, \mathbb{R})^*)$ by $\Phi(v, \theta)(T) = \theta(Tv)$. These maps induce a map

$$(M, \Phi) : \mathcal{F}_*(\mathbb{R}^d) \rightarrow \mathcal{F}_*(\mathfrak{sl}(d, \mathbb{R})).$$

Say that two points $(v, \theta), (w, \varphi) \in \mathcal{F}_*(\mathbb{R}^d)$ are in *general position* if

$$\theta(w) \neq 0 \text{ and } \varphi(v) \neq 0.$$

Lemma 4.6. *The maps M and Φ are Ad-equivariant. If $(v, \theta), (w, \varphi) \in \mathcal{F}_*(\mathbb{R}^d)$ are in general position then the points*

$$(M, \Phi)(v, \theta) \text{ and } (M, \Phi)(w, \varphi)$$

are also in general position. If g and g^{-1} are proximal then $\text{Ad } g$ is proximal and its attractor $(\text{Ad } g)_+$ is $M(g_+, (g^{-1})_-)$.

The proof of the Lemma is standard and direct.

Lemma 4.7. *Consider a convex irreducible representation $\rho : \Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ and consider the map $\eta = M \circ (\xi, \xi^*) : \partial_\infty \Gamma \rightarrow \mathbb{P}(\mathfrak{sl}(d, \mathbb{R}))$. Denote by $V = \langle \eta(\partial_\infty \Gamma) \rangle$ and*

$$\eta^* = (\Phi \circ (\xi, \xi^*)) \cap V.$$

Then $A_\rho = \text{Ad} \circ \rho|V : \Gamma \rightarrow \text{PGL}(V)$ is an irreducible convex representation with equivariant maps (η, η^*) , moreover for a non torsion $\gamma \in \Gamma$ then

$$\lambda_1(A_\rho \gamma) = \lambda_1(\rho \gamma) - \lambda_d(\rho \gamma).$$

We will say that A_ρ is the *irreducible adjoint representation* of ρ .

Proof. Irreducibility follows from Lemma 4.2 The other properties are consequence of Lemma 4.6 together with Lemma 4.3. The last statement follows from the fact that if $\gamma \in \Gamma$ is non torsion then $\xi^*(\gamma_+)$ is the repelling hyperplane of $\rho\gamma^{-1}$ and the attractor of A_ρ ,

$$M(\xi(\gamma_+), \xi^*(\gamma_+)),$$

belongs to V . □

Remark 4.8. Remark that the Hilbert entropy verifies $H_\rho = 2h_{A_\rho} < \infty$.

Hence, using Theorem 4.5 one obtains:

Corollary 4.9. *Let $\rho : \Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ be an irreducible convex representation, then there exists a positive Hölder-continuous function $f : \text{U}\Gamma \rightarrow \mathbb{R}_+^*$ such that*

$$\int_{[\gamma]} f = \frac{\lambda_1(\rho \gamma) - \lambda_d(\rho \gamma)}{2}$$

for every non torsion $[\gamma]$.

4.2. Regularity. The following Lemma is from Benoist [4].

Lemma 4.10 (Benoist [4]). *Let $g \in \text{PGL}(V)$ be proximal and let $V_{\lambda_2(g)}$ be the sum of the characteristic spaces of g whose eigenvalue is of modulus $\exp \lambda_2(g)$. Then for every $v \notin \mathbb{P}(g_-)$ with non zero component in $V_{\lambda_2(g)}$ one has*

$$\lim_{n \rightarrow \infty} \frac{\log d_{\mathbb{P}}(g^n(v), g_+)}{n} = \lambda_2(g) - \lambda_1(g).$$

The following Lemma is the key to relate the Hölder exponent of the equivariant map of a convex representation and eigenvalues of $\rho(\gamma)$ for non torsion $\gamma \in \Gamma$.

Lemma 4.11. *Let $\rho : \Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ be a convex irreducible representation and denote by α the Hölder exponent of the equivariant mapping ξ , then for every non torsion $\gamma \in \Gamma$ one has*

$$\alpha \leq \min \left\{ \frac{\lambda_1(\rho \gamma) - \lambda_2(\rho \gamma)}{|\gamma|}, \frac{\lambda_{d-1}(\rho \gamma) - \lambda_d(\rho \gamma)}{|\gamma|} \right\}.$$

Proof. Consider a non torsion $\gamma \in \Gamma$. Since ρ is irreducible there exists $x \in \partial_\infty \Gamma - \{\gamma_-\}$ such that $\xi(x)$ has non zero projection to $V_{\lambda_2(\rho \gamma)}$, the characteristic space of $\rho \gamma$ of eigenvalue of modulus $\exp \lambda_2(\rho \gamma)$. Lemma 4.3 states that $\xi(\gamma_+)$ is the attractor of $\rho \gamma$, hence applying Benoist's Lemma 4.10 we obtain

$$\lambda_2(\rho \gamma) - \lambda_1(\rho \gamma) = \lim_{n \rightarrow \infty} \frac{\log d_{\mathbb{P}}(\rho \gamma^n(\xi x), \xi(\gamma_+))}{n} = \lim_{n \rightarrow \infty} \frac{\log d_{\mathbb{P}}(\xi(\gamma^n x), \xi(\gamma_+))}{n}$$

since ξ is equivariant. Hölder continuity of ξ implies that the last quantity is smaller than

$$\lim_{n \rightarrow \infty} \frac{\log K \delta_o(\gamma^n x, \gamma_+)^{\alpha}}{n} = -\alpha |\gamma|$$

according to Lemma 3.1. Thus, for every $\gamma \in \Gamma$, one has

$$\alpha \leq \frac{\lambda_1(\rho\gamma) - \lambda_2(\rho\gamma)}{|\gamma|},$$

applying this inequality to γ^{-1} one obtains

$$\alpha \leq \frac{\lambda_{d-1}(\rho\gamma) - \lambda_d(\rho\gamma)}{|\gamma|}.$$

□

5. PROOF OF THEOREM A

This section is devoted to the proof of Theorem A. Consider an irreducible convex representation $\rho : \Gamma \rightarrow \mathrm{PGL}(d, \mathbb{R})$ and denote by c either the Hölder cocycle

$$\beta_\rho \text{ or } \frac{\beta_\rho + \bar{\beta}_\rho}{2},$$

and denote by $\ell_c(\gamma)$ the period of a non torsion γ for c . This is to say, either $\ell_c(\gamma)$ equals $\lambda_1(\rho\gamma)$ or either it equals $(\lambda_1(\rho\gamma) - \lambda_d(\rho\gamma))/2$.

Recall that

$$h_c = \lim_{t \rightarrow \infty} \frac{\log \#\{[\gamma] \in [\Gamma] \text{ non torsion} : \ell_c(\gamma) \leq t\}}{t}.$$

According to Corollary 4.5 or to Corollary 4.9 (depending on whether $c = \beta_\rho$ or $c = (\beta_\rho + \bar{\beta}_\rho)/2$) there exists a positive Hölder-continuous function $f : \mathrm{U}\Gamma \rightarrow \mathbb{R}_+^*$ such that for every non torsion conjugacy class $[\gamma]$ of $[\Gamma]$ one has

$$\int_{[\gamma]} f = \ell_c(\gamma).$$

Let $m_{-h_c f}$ denote the equilibrium state of $-h_c f$ and consider a sequence of periodic orbits $\{[\gamma_n]\}$ such that

$$\frac{\mathrm{Leb}_{\gamma_n}}{|\gamma_n|} \rightarrow m_{-h_c f}$$

as $n \rightarrow \infty$. The existence of this sequence is guaranteed by ergodicity of equilibrium states (Proposition 2.6) and Anosov's closing Lemma 2.5.

One thus obtains

$$\frac{\ell_c(\gamma_n)}{|\gamma_n|} = \frac{1}{|\gamma_n|} \int_{[\gamma_n]} f \rightarrow \int f dm_{-h_c f}$$

which, using Lemma 2.7, is equal to

$$\frac{h(\phi, m_{-h_c f})}{h_c}.$$

Hence, given $\varepsilon > 0$ one has

$$\frac{\ell_c(\gamma_n)}{|\gamma_n|} \leq \frac{h(\phi, m_{-h_c f})}{h_c} (1 + \varepsilon)$$

for all n large enough. Using Lemma 4.11 one has

$$\begin{aligned} \alpha &\leq \min \left\{ \frac{(\lambda_1 - \lambda_2)(\rho\gamma_n)}{|\gamma_n|}, \frac{(\lambda_{d-1} - \lambda_d)(\rho\gamma_n)}{|\gamma_n|} \right\} \\ &\leq \min \left\{ \frac{(\lambda_1 - \lambda_2)(\rho\gamma_n)}{\ell_c(\gamma)}, \frac{(\lambda_{d-1} - \lambda_d)(\rho\gamma_n)}{\ell_c(\gamma)} \right\} \frac{h(\phi, m_{-h_c f})}{h_c} (1 + \varepsilon). \end{aligned}$$

Theorem 3.3 states that h_Γ is the maximal entropy of the flow ϕ on $\mathbf{U}\Gamma$, hence

$$\frac{\alpha h_c}{h_\Gamma} \leq \frac{\alpha h_c}{h(\phi, m_{-h_c f})} \leq \min \left\{ \frac{(\lambda_1 - \lambda_2)(\rho\gamma_n)}{\ell_c(\gamma)}, \frac{(\lambda_{d-1} - \lambda_d)(\rho\gamma_n)}{\ell_c(\gamma)} \right\} (1 + \varepsilon) \quad (7)$$

for all n large enough.

We will now distinguish the two cases:

First case: $c = \beta_\rho$. In this case $\ell_c(\gamma) = \lambda_1(\rho\gamma)$, $h_c = h_\rho$ (the spectral entropy of ρ) and equation (7) is

$$\frac{\alpha h_\rho}{h_\Gamma} \leq \frac{\alpha h_\rho}{h(\phi, m_{-h_\rho f})} \leq \min \left\{ \frac{\lambda_1 - \lambda_2}{\lambda_1}(\rho\gamma_n), \frac{\lambda_{d-1} - \lambda_d}{\lambda_1}(\rho\gamma_n) \right\} (1 + \varepsilon).$$

We will now maximize the function $V_1 : \mathbb{P}(\mathfrak{a}^+) \rightarrow \mathbb{R}$ defined by

$$V_1(a_1, \dots, a_d) = \min \left\{ \frac{a_1 - a_2}{a_1}, \frac{a_{d-1} - a_d}{a_1} \right\}.$$

Recall that

$$\mathfrak{a}^+ = \{(a_1, \dots, a_d) \in \mathbb{R}^d : a_1 + \dots + a_d = 0 \text{ and } a_1 \geq \dots \geq a_d\}$$

and consider $a \in \mathfrak{a}^+$. We will distinguish two cases.

Assume $a_2 \geq 0$: In this case one has

$$V_1(a) \leq \frac{a_1 - a_2}{a_1} = 1 - \frac{a_2}{a_1} \leq 1.$$

Assume $a_2 < 0$:

Remark 5.1. In this case one has $a_1 - a_2 > a_{d-1} - a_d$, hence $V_1(a) = (a_{d-1} - a_d)/a_1$.

Proof. Remark that $a_{k+1} - a_k \leq 0$ for all $k \in \{1, \dots, d-1\}$. Using the following tricky equality (recall $d \geq 3$)

$$a_1 + (d-1)a_2 + \sum_{k=2}^{d-1} (d-k)(a_{k+1} - a_k) = a_1 + a_2 + \dots + a_d = 0$$

one obtains

$$a_1 - a_2 + a_d - a_{d-1} = -da_2 - \sum_{k=2}^{d-2} (d-k)(a_{k+1} - a_k) > 0.$$

Hence $a_1 - a_2 > a_{d-1} - a_d$.

□

Since $0 > a_2 \geq \dots \geq a_d$ one has

$$a_1 = -a_2 - \dots - a_d > -(a_{d-1} + a_d) \geq 0.$$

Given that $d \geq 3$ one obtains, $a_{d-1} < 0 < -a_{d-1}$ and subtracting a_d on each side one gets $a_{d-1} - a_d < -(a_{d-1} + a_d) < a_1$, finally

$$V_1(a) = \frac{a_{d-1} - a_d}{a_1} < 1.$$

In any case one obtains $V_1 \leq 1$. We then get

$$\frac{\alpha h_\rho}{h_\Gamma} \leq \frac{\alpha h_\rho}{h(\phi, m_{-h_\rho f})} \leq V_1(\lambda(\rho\gamma_n))(1 + \varepsilon) \leq 1 + \varepsilon, \quad (8)$$

since ε is arbitrary we obtain the desired inequality.

Remark 5.2. Consider

$$\alpha_\rho = \sup\{\alpha \in (0, 1] : \xi \text{ is } \alpha\text{-H\"older}\}.$$

We remark that if $\alpha_\rho h_\rho = h_\Gamma$ then inequality (8) implies that $m_{-h_\rho f}$ is the measure of maximal entropy of ϕ and thus (using Proposition 2.6) the function $f : \mathcal{U}\Gamma \rightarrow \mathbb{R}_+^*$ is Livšic-cohomologous to a constant, i.e. there exists $c > 0$ such that

$$\lambda_1(\rho\gamma) = c|\gamma|$$

for every non torsion $\gamma \in \Gamma$.

Second case: $c = (\beta_\rho + \bar{\beta}_\rho)/2$. In this case we have $\ell_c(\gamma) = (\lambda_1(\rho\gamma) - \lambda_d(\rho\gamma))/2$, $h_c = H_\rho$ (the Hilbert entropy of ρ) and inequality (7) states

$$\frac{\alpha H_\rho}{h_\Gamma} \leq \frac{\alpha H_\rho}{h(\phi, m_{-H_\rho f})} \leq \min \left\{ \frac{\lambda_1 - \lambda_2}{(\lambda_1 - \lambda_d)/2}(\rho\gamma_n), \frac{\lambda_{d-1} - \lambda_d}{(\lambda_1 - \lambda_d)/2}(\rho\gamma_n) \right\} (1 + \varepsilon)$$

for all n large enough.

We will now maximize the function $V_2 : \mathbb{P}(\mathfrak{a}^+) \rightarrow \mathbb{R}$ defined by

$$V_2(a_1, \dots, a_d) = \min \left\{ \frac{a_1 - a_2}{(a_1 - a_d)/2}, \frac{a_{d-1} - a_d}{(a_1 - a_d)/2} \right\}.$$

Consider $a \in \mathfrak{a}^+$ such that

$$x := a_1 - a_2 \leq a_{d-1} - a_d =: y.$$

For such a one has $a_2 = a_1 - x$ and $a_{d-1} = y + a_d$. Since $d \geq 3$ one has $a_2 \geq a_{d-1}$ hence $a_1 - x \geq a_d + y \geq a_d + x$ and thus

$$V_2(a) = \frac{2x}{a_1 - a_d} \leq 1.$$

If, on the opposite, one has $a \in \mathfrak{a}^+$ such that

$$x := a_{d-1} - a_d \leq a_1 - a_2 =: y,$$

then, again the fact that $a_2 \geq a_{d-1}$ implies $a_1 - x \geq a_1 - y \geq a_d + x$ and thus

$$V_2(a) = \frac{2x}{a_1 - a_d} \leq 1.$$

In any case one obtains $V_2 \leq 1$. We then get

$$\frac{\alpha H_\rho}{h_\Gamma} \leq \frac{\alpha H_\rho}{h(\phi, m_{-H_\rho f})} \leq V_2(\lambda(\rho\gamma_n))(1 + \varepsilon) \leq 1 + \varepsilon, \quad (9)$$

since ε is arbitrary we obtain the desired inequality.

Remark 5.3. Remark that if $\alpha_\rho H_\rho = h_\Gamma$ then inequality (9) implies that $m_{-H_\rho f}$ is the measure of maximal entropy of ϕ and thus (using Proposition 2.6) the function $f : \mathcal{U}\Gamma \rightarrow \mathbb{R}_+^*$ is Livšic-cohomologous to a constant, i.e. there exists $c > 0$ such that

$$\lambda_1(\rho\gamma) - \lambda_d(\rho\gamma) = 2c|\gamma|$$

for every non torsion $\gamma \in \Gamma$.

6. CROSS RATIOS: GENERAL LEMMAS AND A THEOREM OF LABOURIE

Consider a compact metric space X and

$$X^{(4)} = \{(x, y, z, t) \in X^4 \text{ such that } x \neq t \text{ and } y \neq z\}$$

Definition 6.1. A cross ratio on X is a Hölder-continuous function $\mathbf{b} : X^{(4)} \rightarrow \mathbb{R}$ that verifies the following relations for every $x, y, z, t, w \in X$:

- $\mathbf{b}(x, y, z, t) = \mathbf{b}(z, t, x, y)$,
- $\mathbf{b}(x, y, x, t) = 1 = \mathbf{b}(x, y, z, y)$,[†]
- $\mathbf{b}(x, y, z, t) = 0$ if and only if $x = y$ or $z = t$,
- $\mathbf{b}(x, y, z, t) = \mathbf{b}(x, y, z, w)\mathbf{b}(x, w, z, t)$,
- $\mathbf{b}(x, y, z, t) = \mathbf{b}(x, y, w, t)\mathbf{b}(w, y, z, t)$.

Remark that if \mathbf{b} is a cross ratio then so are \mathbf{b}^n for $n \in \mathbb{Z}$, its modulus $|\mathbf{b}|$, and $|\mathbf{b}|^c$ for any positive real number c .

The following Lemma is trivial.

Lemma 6.2. *Consider a cross ratio \mathbf{b} on X and assume a connected group G acts on X , then for every $g \in G$ and $(x, y, z, t) \in X^{(4)}$ the sign of $\mathbf{b}(x, y, z, t)$ and the sign of $\mathbf{b}(gx, gy, gz, gt)$ coincide.*

Remark that if for every pair of points $x, y \in X$ the space $X - \{x, y\}$ is connected then any cross ratio \mathbf{b} defined on X is necessarily non negative. Indeed, fixing $x, y, z \in X$ pairwise distinct and considering $\mathbf{b}(x, \cdot, y, z) : X - \{y\} \rightarrow \mathbb{R}$ one obtains a function that only vanishes on x , i.e. $\mathbf{b}(x, \cdot, y, z) : X - \{x, y\} \rightarrow \mathbb{R}$ is continuous, non vanishing and takes the value 1 on z . Analog reasoning works for the other entries of \mathbf{b} .

Lemma 6.3. *Let Γ be a hyperbolic group that does not split over a cyclic \mathbb{Z} (recall definition 1.8), then any cross ratio defined on $\partial_\infty \Gamma$ is non negative.*

Proof. This is direct consequence of the preceding paragraph and following deep fact of Bowditch [8]: the boundary of such group does not have local cut points on $\partial_\infty \Gamma$, which in turn implies that $\partial_\infty \Gamma - \{x, y\}$ is connected for any pair of points on $x, y \in \partial_\infty \Gamma$. \square

The rank of a cross ratio is defined as follows. Consider X^{p*} the set of pairs $(e, u) = (e_0, \dots, e_p, u_0, \dots, u_p)$ of $p + 1$ -tuples on X such that $e_j \neq e_i \neq u_0$ and $u_j \neq u_i \neq e_0$ when $j > i > 0$, define then

$$\chi_{\mathbf{b}}^p(e, u) = \det_{i, j > 0} (\mathbf{b}(e_i, u_j, e_0, u_0)).$$

Definition 6.4. The rank^\dagger of a cross ratio \mathbf{b} on X is defined by

$$\text{rank } \mathbf{b} = \inf\{p \in \mathbb{N} : \chi_{\mathbf{b}}^p \equiv 0\} - 1,$$

if it exists.

Consider $d \in \mathbb{N}$, a *convex map* from X to $\mathbb{P}(\mathbb{R}^d)$ is a Hölder-continuous map

$$(\xi, \xi^*) : X \rightarrow \mathbb{P}(\mathbb{R}^d) \times \mathbb{P}(\mathbb{R}^{d*})$$

such that $\xi(x) \subset \ker \xi^*(y)$ if and only if $x = y$.

[†]This condition is weaker than the analog one defined on Labourie [17].

[‡]This is a slightly different definition of rank than the one from Labourie [18].

Given a convex map from X to $\mathbb{P}(\mathbb{R}^d)$ one can define a cross ratio on X $\mathbf{b}^\xi : X^{(4)} \rightarrow \mathbb{R}$ by

$$\mathbf{b}^\xi(x, y, z, t) = \frac{\varphi(u) \psi(v)}{\psi(u) \varphi(v)} \quad (10)$$

where $\varphi \in \xi^*(x)$, $\psi \in \xi^*(z)$, $u \in \xi(y)$ and $v \in \xi(t)$. Remark that the result does not depend on the choice of φ , ψ , u and v made.

Remark 6.5 (Labourie [18]). If $\xi(X)$ generates \mathbb{R}^d then \mathbf{b}^ξ has rank d .

It turns out that the converse is also true in a stronger version.

Theorem 6.6 (Labourie [18]). *Consider a rank d cross ratio $\mathbf{b} : X^{(4)} \rightarrow \mathbb{R}$ and assume a group G acts on X leaving \mathbf{b} invariant, then there exists a representation $\rho : G \rightarrow \mathrm{PGL}(d, \mathbb{R})$ and a ρ -equivariant convex map*

$$(\xi, \xi^*) : X \rightarrow \mathbb{P}(\mathbb{R}^d) \times \mathbb{P}(\mathbb{R}^{d*})$$

such that $\xi(X)$ generates \mathbb{R}^d and $\mathbf{b} = \mathbf{b}^\xi$. The representation and the convex map are unique modulo conjugation via $\mathrm{PGL}(d, \mathbb{R})$.

Labourie's original statement is for $X = S^1$ the circle, and $G = \pi_1 \Sigma$ the fundamental group of a closed hyperbolic surface, nevertheless the proof of existence is an exact copy of Proposition 5.7 of Labourie [18] and the proof of uniqueness is an exact copy of Lemma 4.3 of Labourie [18].

The main consequence we are interested in is the existence of a linear action of G on \mathbb{R}^d when G acts on X preserving a rank d cross ratio.

We will need the following Lemma for the proof of Hilbert's entropy rigidity.

Lemma 6.7. *Consider a convex map $(\xi, \xi^*) : X \rightarrow \mathbb{P}(\mathbb{R}^d) \times \mathbb{P}(\mathbb{R}^{d*})$ and the adjoint equivariant maps*

$$(\eta, \eta^*) := (M(\xi, \xi^*), \Phi(\xi, \xi^*)) : X \rightarrow \mathbb{P}(\mathfrak{sl}(d, \mathbb{R})) \times \mathbb{P}(\mathfrak{sl}(d, \mathbb{R})^*),$$

defined on Section 4. Then

$$\mathbf{b}^\eta(x, y, z, t) = \mathbf{b}^\xi(x, y, z, t) \mathbf{b}^\xi(y, x, t, z).$$

Proof. The proof is an explicit calculation. \square

7. THE CROSS RATIO OF A HÖLDER COCYCLE

Consider now Γ a convex co-compact group of a $\mathrm{CAT}(-1)$ space X . Two Hölder cocycles $c, c' : \Gamma \times \partial_\infty \Gamma \rightarrow \mathbb{R}$ are said to be *cohomologous* if there exists a Hölder-continuous function $U : \partial_\infty \Gamma \rightarrow \mathbb{R}$ such that for all $\gamma \in \Gamma$ one has

$$c(\gamma, x) - c'(\gamma, x) = U(\gamma x) - U(x).$$

One easily deduces from the definition that the set of periods $\{\ell_c(\gamma) : \gamma \in \Gamma\}$ of a Hölder cocycle is a cohomological invariant.

Theorem 7.1 (Ledrappier [19]). *Two Hölder cocycles are cohomologous if and only if they have the same period for every non torsion $\gamma \in \Gamma$.*

We shall be interested in cocycles whose periods are positive, i.e. such that $\ell_c(\gamma) > 0$ for every non torsion $\gamma \in \Gamma$. The *entropy*[†] of such cocycle is defined by:

$$h_c = \limsup_{t \rightarrow \infty} \frac{1}{t} \log \# \{[\gamma] : \ell_c(\gamma) \leq t\} \in \mathbb{R}_+ \cup \{\infty\}.$$

[†]In [20] this is called the exponential growth rate of the cocycle.

Given a Hölder cocycle there exists a *dual Hölder cocycle* \bar{c} determined by the equation

$$\ell_{\bar{c}}(\gamma) = \ell_c(\gamma^{-1})$$

(see Ledrappier [19] or [20] for details). Denote by

$$\partial_{\infty}^{(2)}\Gamma = \partial_{\infty}\Gamma \times \partial_{\infty}\Gamma - \{(x, x) : x \in \partial_{\infty}\Gamma\}.$$

A *Gromov product* for the ordered pair (c, \bar{c}) is a function $[\cdot, \cdot] = [\cdot, \cdot]_{(c, \bar{c})} : \partial_{\infty}^{(2)}\Gamma \rightarrow \mathbb{R}$ such that for every $\gamma \in \Gamma$ and $x, y \in \partial_{\infty}\Gamma$ distinct one has

$$[\gamma x, \gamma y] - [x, y] = -(\bar{c}(\gamma, x) + c(\gamma, y))^{\dagger}.$$

Theorem 7.2 (Ledrappier [19]). *Given a Hölder cocycle c there exists a dual cocycle \bar{c} . If c has finite and positive entropy then there exists a Gromov product for the pair (c, \bar{c}) . Moreover, if $y \rightarrow x$ then $[x, y] \rightarrow -\infty$.*

We will hence adopt the convention $[x, x] = -\infty$.

Lemma 7.3. *Let $\{c, \bar{c}\}$ be a pair of dual Hölder cocycles with positive and finite entropy, and $[\cdot, \cdot]$ a Gromov product for the ordered pair (c, \bar{c}) . Then the function $\mathbf{b}_c : \partial_{\infty}\Gamma^{(4)} \rightarrow \mathbb{R}$ defined by*

$$\mathbf{b}_c(x, y, z, t) = \exp\{[x, y] - [z, y] + [z, t] - [x, t]\},$$

is a Γ -invariant cross ratio on $\partial_{\infty}\Gamma$. Moreover, \mathbf{b}_c only depends on the cohomology class of c . In particular the cross ratio \mathbf{b}_c is uniquely determined by the periods $\{\ell_c(\gamma) : \gamma \in \Gamma\}$ of c .

Hence, we define the *cross ratio*[§] associated to a Hölder cocycle c as the function \mathbf{b}_c of the last Lemma. Remark that by definition $\mathbf{b}_c \geq 0$.

Proof. The properties from the definition 6.1 are easily verified and Γ -invariance follows from the definition of Gromov product.

Let us focus on the fact that \mathbf{b}_c only depends on the cohomology class of c . Remark that Ledrappier's Theorem 7.1 implies that the cohomology class of \bar{c} is determined by the cohomology class of c . Consider then c' a cocycle cohomologous to c , i.e. $c'(\gamma, x) - c(\gamma, x) = U(\gamma x) - U(x)$. The Gromov product associated to the ordered pair (c', \bar{c}) is $[x, y]' = [x, y] + U(y)$. One then has

$$[x, y]' - [z, y]' = [x, y] + U(y) - [z, y] - U(y) = [x, y] - [z, y].$$

Hence, the cross ratios associated to (c, \bar{c}) and (c', \bar{c}) coincide. analog reasoning works for a cocycle cohomologous to \bar{c} . The last statement follows from this and Ledrappier's Theorem 7.1. \square

It would be nice to have Benoist-like formula (see Benoist [3, Lemma 1.6]):

Question. Let c be a Hölder cocycle with finite and positive entropy, is it true that for every non torsion $\gamma, h \in \Gamma$ one has

$$\exp\{\ell_c(\gamma^n h^n) - \ell_c(\gamma^n) - \ell_c(h^n)\} \rightarrow \mathbf{b}_c(\gamma_-, h_+, h_-, \gamma_+)$$

as $n \rightarrow \infty$?

[†]Remark this definition applied to a CAT(−1)-space gives the opposite of the usual Gromov product.

[§]This is Ledrappier's [19] construction of a cross ratio, but in our case the Hölder cocycle is not necessarily what he calls pair.

The following Lemma clarifies the relation between the cross ratios \mathbf{b}_c and $\mathbf{b}_{\bar{c}}$. I would like to thank Qiongling Li for discussions concerning this matter.

Lemma 7.4. *Consider a Hölder cocycle c with finite and positive entropy. Then for every $(x, y, z, t) \in \partial_\infty \Gamma^{(4)}$ one has*

$$\mathbf{b}_c(x, y, z, t) = \mathbf{b}_{\bar{c}}(y, x, t, z)$$

and

$$\mathbf{b}_{c+\bar{c}}(x, y, z, t) = \mathbf{b}_c(x, y, z, t) \mathbf{b}_{\bar{c}}(x, y, z, t).$$

Proof. Let $[\cdot, \cdot]_1$ be the Gromov product for the ordered pair (c, \bar{c}) . An explicit computation shows that $[\cdot, \cdot]_2 : \partial_\infty^{(2)} \Gamma \rightarrow \mathbb{R}$ defined by

$$[x, y]_2 = [y, x]_1$$

is a Gromov product for the ordered pair (\bar{c}, c) . Hence the first equality in the Lemma. The second equality follows then by remarking that $[x, y]_3 = [x, y]_1 + [x, y]_2$ is a Gromov product for $(c + \bar{c}, c + \bar{c})$. \square

8. CONVEX REPRESENTATIONS AND CROSS RATIOS

Consider now a convex representation $\rho : \Gamma \rightarrow \mathrm{PGL}(d, \mathbb{R})$ with equivariant maps $\xi : \partial_\infty \Gamma \rightarrow \mathbb{P}(\mathbb{R}^d)$ and $\xi^* : \partial_\infty \Gamma \rightarrow \mathbb{P}(\mathbb{R}^{d*})$. Lemma 4.3 implies that (ξ, ξ^*) is a convex map in the sense of Section 6, we define then the *cross ratio of ρ* to be the cross ratio of the equivariant map ξ , i.e.

$$\mathbf{b}_\rho = \mathbf{b}^\xi.$$

Consider now the Hölder cocycles $\beta_\rho, \bar{\beta}_\rho : \Gamma \times \partial_\infty \Gamma \rightarrow \mathbb{R}$ defined on Section 4. Lemma 4.4 implies that $\{\beta_\rho, \bar{\beta}_\rho\}$ is a pair of dual cocycles.

Consider now the function $\mathcal{G} : \mathbb{P}(\mathbb{R}^{d*}) \times \mathbb{P}(\mathbb{R}^d) - \{(\varphi, v) : \varphi(v) = 0\} \rightarrow \mathbb{R}$ defined by

$$\mathcal{G}(\theta, v) = \log \frac{|\theta(v)|}{\|\theta\| \|v\|}$$

and define $[\cdot, \cdot] : \partial_\infty^{(2)} \Gamma \rightarrow \mathbb{R}$ by

$$[x, y] = \mathcal{G}(\xi^*(x), \xi(y)).$$

A direct computation shows that the function $[\cdot, \cdot] : \partial_\infty^{(2)} \Gamma \rightarrow \mathbb{R}$ is a Gromov product for the ordered pair $(\beta_\rho, \bar{\beta}_\rho)$. The cross ratio associated to the cocycle β_ρ is

$$\mathbf{b}_{\beta_\rho}(x, y, z, t) = \left| \frac{\varphi(u) \psi(v)}{\psi(u) \varphi(v)} \right|$$

where $\varphi \in \xi^*(x)$, $\psi \in \xi^*(z)$, $u \in \xi(y)$ and $v \in \xi(t)$. Hence $\mathbf{b}_{\beta_\rho} = |\mathbf{b}_\rho|$. Remark also that the cross ratio associated to \mathbf{A}_ρ , the irreducible adjoint representation of ρ , verifies

$$|\mathbf{b}_{\mathbf{A}_\rho}| = \mathbf{b}_{\beta_\rho + \bar{\beta}_\rho}.$$

As a conclusion of this section we state the following consequence of Lemma 7.3:

Lemma 8.1. *Let $\rho : \Gamma \rightarrow \mathrm{PSL}(d, \mathbb{R})$ be an irreducible convex representation, then the cross ratio $|\mathbf{b}_\rho|$ is induced by the Hölder cocycle β_ρ , and is thus uniquely determined by $\{\lambda_1(\rho\gamma) : \gamma \in \Gamma\}$. Moreover the cross ratio $|\mathbf{b}_{\mathbf{A}_\rho}|$ is induced by the Hölder cocycle $\beta_\rho + \bar{\beta}_\rho$ and is thus uniquely determined by $\{\lambda_1(\rho\gamma) - \lambda_d(\rho\gamma) : \gamma \in \Gamma\}$.*

9. PROXIMAL REPRESENTATIONS OF SEMI-SIMPLE LIE GROUPS

Consider a real algebraic non compact semi-simple Lie group G . Let K be a maximal compact subgroup of G and τ the Cartan involution on \mathfrak{g} whose fixed point set is the Lie algebra of K . Consider $\mathfrak{p} = \{v \in \mathfrak{g} : \tau v = -v\}$ and \mathfrak{a} a maximal abelian subspace contained in \mathfrak{p} .

Let Σ be the set of (restricted) roots of \mathfrak{a} on \mathfrak{g} , \mathfrak{a}^+ a closed Weyl chamber, Σ^+ a system of positive roots on Σ associated to \mathfrak{a}^+ and denote by Π the set of simple roots associated to the choice Σ^+ .

For $g \in G$ denote by $\lambda(g) \in \mathfrak{a}^+$ its *Jordan projection*, this is the unique element on \mathfrak{a}^+ such that $\exp \lambda(g)$ is conjugated to the \mathbb{R} -regular element on the Jordan decomposition of g .

For an irreducible representation $\phi : G \rightarrow \mathrm{PGL}(d, \mathbb{R})$ denote by $\chi_\phi \in \mathfrak{a}^*$ its *restricted highest weight*. For every $g \in G$ one has, by definition,

$$\lambda_1(\phi g) = \chi_\phi(\lambda(g)). \quad (11)$$

We say that ϕ is *proximal* if $\phi(G)$ contains a proximal element. One has the following standard proposition in Representation Theory.

Proposition 9.1 (see Benoist [5, Section 2.2]). *The set of restricted weights of \mathfrak{a}^* is in bijection with (equivalence classes of) irreducible proximal representations of G .*

Let W be the Weyl group of Σ and denote $u_0 : \mathfrak{a} \rightarrow \mathfrak{a}$ the biggest element in W , u_0 is the unique element in W that sends \mathfrak{a}^+ to $-\mathfrak{a}^+$. The *opposition involution* $i : \mathfrak{a} \rightarrow \mathfrak{a}$ is the defined by $i := -u_0$.

Consider $\{\omega_\theta\}_{\theta \in \Pi}$ the set of fundamental weights of Π . We will need the following result of Tits [23].

Proposition 9.2 (Tits [23]). *For each $\theta \in \Pi$ there exists a finite dimensional proximal irreducible representation $\Lambda_\theta : G \rightarrow \mathrm{PGL}(V_\theta)$ such that the restricted highest weight χ_θ of Λ_θ is an integer multiple of the fundamental weight ω_θ .*

We will now specify on the group $\mathrm{Isom} \mathbb{H}^k$. The Cartan subspace $\mathfrak{a}_{\mathbb{H}^k}$ is 1-dimensional and is thus identified with \mathbb{R} . The *Jordan projection* of $\gamma \in \mathrm{Isom} \mathbb{H}^k$ is

$$\lambda_{\mathbb{H}^k}(\gamma) = \inf_{p \in \mathbb{H}^k} d_{\mathbb{H}^k}(p, \gamma p),$$

and thus $\lambda_{\mathbb{H}^k}(\gamma)$ coincides with the translation length $|\gamma|$ when γ is a hyperbolic element.

Remark 9.3. Remark that if $\rho : \mathrm{Isom} \mathbb{H}^k \rightarrow \mathrm{PGL}(k+1, \mathbb{R})$ is the Klein model of \mathbb{H}^k and $\gamma \in \mathrm{Isom} \mathbb{H}^k$ is hyperbolic then $\lambda_1(\rho\gamma) = |\gamma|$ and $\lambda_1(\mathrm{Ad} \rho\gamma) = 2|\gamma|$.

10. COMPUTING THE RANK OF $\mathbb{B}_{\mathbb{H}^k}$

Consider $\phi : \mathrm{Isom} \mathbb{H}^k \rightarrow \mathrm{PGL}(k+1, \mathbb{R})$ the Klein model of \mathbb{H}^k . As explained before we have a convex map

$$(\xi, \xi^*) : \partial_\infty \mathbb{H}^k \rightarrow \mathbb{P}(\mathbb{R}^{k+1}) \times \mathbb{P}(\mathbb{R}^{k+1}^*).$$

Define the cross ratio $\mathbb{B}_{\mathbb{H}^k}$ on $\partial_\infty \mathbb{H}^k$ by[†]

$$\mathbb{B}_{\mathbb{H}^k} = \mathbf{b}^\xi.$$

[†]For $k = 2$ this is the square of the usual cross ratio on the circle.

Remark 10.1. Remark that $\mathbb{B}_{\mathbb{H}^k}$ is non negative: this is obvious for $k \geq 3$ and for $k = 2$ one uses the fact that $\xi(\partial_\infty \mathbb{H}^2) \subset \mathbb{P}(\mathbb{R}^3)$ is the boundary of a convex set. Remark also that the rank of $\mathbb{B}_{\mathbb{H}^k}$ is $k + 1$.

As observed before, if α is a positive real number then $\mathbb{B}_{\mathbb{H}^k}^\alpha$ is also a cross ratio.

Remark 10.2. Let Γ be a Zariski dense convex co-compact subgroup of $\text{Isom } \mathbb{H}^k$ and α a positive real number, then the cross ratio $\mathbb{B}_{\mathbb{H}^k}^\alpha|_{L_\Gamma}$ is induced by a Hölder cocycle (namely α -Busseman's cocycle) and hence it is uniquely determined by the periods of α -Busseman, i.e. by the set $\{\alpha|\gamma| : \gamma \in \Gamma\}$.

Deciding whether the cross ratio $\mathbb{B}_{\mathbb{H}^k}^\alpha$ has finite rank can be a difficult task, the actual definition is hard to handle.

The purpose of this section is to show that, in order to compute the rank of $\mathbb{B}_{\mathbb{H}^k}^\alpha$ one only needs to check on the limit set of a Zariski dense subgroup of $\text{Isom } \mathbb{H}^k$. This is a key step in the proofs of Theorems B and D.

Theorem 10.3. *Consider a Zariski dense subgroup Γ of $\text{Isom } \mathbb{H}^k$ and a positive real number α . If the cross ratio $\mathbb{B}_{\mathbb{H}^k}^\alpha|_{L_\Gamma} : L_\Gamma^{(4)} \rightarrow \mathbb{R}$ has finite rank, then*

$$\text{rank } \mathbb{B}_{\mathbb{H}^k}^\alpha = \text{rank } \mathbb{B}_{\mathbb{H}^k}^\alpha|_{L_\Gamma}.$$

10.1. Polynomial functions on the Furstenberg boundary. We will freely use the notations of Section 9. Let P be a minimal parabolic subgroup of G and denote by $\mathcal{F} = G/P$ the *Furstenberg boundary* of the symmetric space of G . Consider Δ a Zariski dense subgroup of G . We have the following Proposition of Benoist [2].

Proposition 10.4 (Benoist [2]). *The action of Δ on \mathcal{F} has a unique minimal closed invariant set.*

This smallest closed invariant set is called *the limit set* of Δ and denoted by L_Δ . We need the following technical Proposition:

Proposition 10.5. *Consider Δ a Zariski dense subgroup of G . Consider also a finite family of polynomial functions $P_i : \mathcal{F} \rightarrow \mathbb{R}$ for $i \in \{1, \dots, k\}$ and positive real numbers α_i . Consider*

$$f := \sum_1^k \varepsilon_i |P_i|^{\alpha_i} : \mathcal{F} \rightarrow \mathbb{R}$$

where $\varepsilon_i \in \{1, -1\}$ and assume that f vanishes on L_Δ , then f vanishes on \mathcal{F} .

The proof of the Proposition goes by adapting arguments from Section 7 of Benoist [2]. We reproduce here these arguments for completeness. We would like to thank Yves Benoist for discussions that lead to this proof.

We need the following Lemmas from Benoist [2].

Lemma 10.6 (Benoist [2, Lemma 7.1]). *Consider a finite family of 1-parameter subgroups $\{g_i^t : t \in \mathbb{R}\}$ $i \in \{1, \dots, s\}$ of G . Consider the semi-group H generated by*

$$\{g_i^t : i \in \{1, \dots, s\} \text{ and } t \in [1, \infty)\}.$$

If H is Zariski dense in G then H has non empty interior.

An element $g \in G$ is called *semi-simple* if the unipotent element in its Jordan decomposition is trivial.

Lemma 10.7 (Benoist [2]). *Let Δ be a Zariski dense subgroup of G . Then there exists a subgroup Δ' of Δ such that*

- Δ' is still Zariski dense on G ,
- every element of Δ' is semi-simple.

10.2. Hardy fields. Consider \mathcal{K} the ring of germs at infinity of \mathbb{C}^∞ functions defined on a half line $[t_0, \infty)$. A *Hardy field* is a sub-field of \mathcal{K} stable under derivation. Our interest in considering Hardy fields is that if f belongs to a Hardy field then $1/f$ is well defined and hence f has no zeros on some half line $[t, \infty)$.

Proposition 10.8 (Benoist [2]). *Let \mathcal{H} be a Hardy field. Let α be a positive real number and $f \in \mathcal{H}$ then there exists a Hardy field \mathcal{H}' that contains \mathcal{H} and $|f|^\alpha$.*

Recall that G is a real algebraic semi-simple Lie group and that \mathcal{F} is its Furstenberg boundary. Consider a finite family of polynomial functions $P_i : \mathcal{F} \rightarrow \mathbb{R}$ for $i \in \{1, \dots, k\}$ and positive real numbers α_i . Consider

$$f = \sum_{i=1}^k \varepsilon_i |P_i|^{\alpha_i} : \mathcal{F} \rightarrow \mathbb{R}$$

where $\varepsilon_i \in \{1, -1\}$

Lemma 10.9. *Consider the 1-parameter group $\{g^t : t \in \mathbb{R}\}$ generated by an \mathbb{R} -regular element $g \in G$ and fix $x_0 \in \mathcal{F}$. Consider also $h_1, h_2 \in G$. Then the function*

$$t \mapsto f(h_2 g^t h_1 \cdot x_0)$$

either has finite zeros or identically vanishes on some half line.

Proof. For each polynomial P_i , the function

$$t \mapsto P_i(h_2 g^t h_1 \cdot x_0)$$

is a linear combination of functions of the form $t \mapsto e^{kt}$ for some $k \in \mathbb{R}$ (this is consequence of g being \mathbb{R} -regular). All these functions belong to a Hardy field, hence using Proposition 10.8, so does $t \mapsto |P_i(h_2 g^t h_1 \cdot x_0)|^{\alpha_i}$ and thus so does $t \mapsto f(h_2 g^t h_1 \cdot x_0)$. \square

10.3. Proof of Proposition 10.5. We need the following Lemma.

Lemma 10.10 (Benoist [2, Lemma 7.3]). *Let V be a real analytic manifold, $\psi : V \times \mathbb{R} \rightarrow \mathbb{R}$ an analytic function. Consider $Z = \psi^{-1}(0)$ and let $p : Z \rightarrow V$ be the projection on the first coordinate. Assume that $p^{-1}(m)$ is finite for all $m \in V$, then there exists a non empty open set U of V such that $p^{-1}(\overline{U})$ is compact on Z .*

Proof of Proposition 10.5. Fix a point $x_0 \in L_\Delta$ and consider the function $\Phi : G \rightarrow \mathbb{R}$ defined by

$$\Phi(g) = f(g \cdot x_0).$$

Remark that since L_Δ is Δ -invariant and f vanishes on L_Δ one has $\Phi(\Delta) \equiv 0$.

Consider the subgroup Δ' of semi-simple elements of Δ given by Lemma 10.7. Consider H the semi-group generated by $\{g^t : g \in \Delta' \theta \geq 1\}$. Lemma 10.6 implies that H has non empty interior. We will show that Φ vanishes on H , and hence, since Φ is analytic, it will be constant equal to zero.

An element in H is of the form $h_1 \cdots h_s$ where, for each i , one has $h_i = g_i^{t_i}$ for some $g_i \in \Delta'$ and $t_i \in [1, \infty)$. To show that Φ vanishes on H we will proceed by induction, i.e. we will show that if $h_1, h_2 \in H$ and $g \in \Delta'$ are such that

$\Phi(h_2 g^n h_1) = 0$ for every $n \in \mathbb{N}$ then $\Phi(h_2 g^t h_1) = 0$ for all $t \in [1, \infty)$, since $\Phi(\Delta') = 0$ the result will follow.

For $g \in \Delta'$ write $g = m_g a_g$ where m_g is elliptic and a_g is \mathbb{R} -regular and consider the one parameter subgroup $g^t = m_g^t a_g^t$. Let M be the closure of the group

$$\{m_g^t : t \in \mathbb{R}\}.$$

Remark that M is a compact group and hence it is Zariski closed. Consider then

$$Z = \{(m, t) \in M \times [1, \infty) : \Phi(h_2 m a_g^t h_1) = 0\}$$

and $p : Z \rightarrow M$ the projection on the first coordinate.

Remark that for every $m \in M$ one has that $p^{-1}(m)$ is either $m \times [1, \infty)$ or finite. Indeed, Lemma 10.9 states that $t \mapsto \Phi(h_2 m a_g^t h_1)$ belongs to a Hardy field, hence it has finite zeros or it is constant equal to zero.

Assume by contradiction that $Z \neq M \times [1, \infty)$, then there is an open set V of M such that $p^{-1}(m_0)$ is finite for every $m_0 \in V$. Applying Lemma 10.10 implies the existence of an open subset $U \subset V$ such that $p^{-1}(\overline{U})$ is compact, but the set

$$\{n \in \mathbb{N} : m_g^n \in U\}$$

is infinite and, by hypothesis $\Phi(h_2 m_g^n a_g^n h_1) = 0$, hence $(m_g^n, n) \in p^{-1}(\overline{U})$. This contradicts the fact that $p^{-1}(\overline{U})$ is compact. This finishes the proof. \square

10.4. Proof of Theorem 10.3. Let $p_0 \in \mathbb{N}$ be the rank of $\mathbb{B}_{\mathbb{H}^k}^\alpha |_{L_\Gamma}$, this is to say, for every $p > p_0$ the function

$$\chi^p(e, u) := \chi_{\mathbb{B}_{\mathbb{H}^k}^\alpha |_{L_\Gamma}}^p(e, u) = \det_{i,j>0} (\mathbb{B}_{\mathbb{H}^k}(e_i, u_j, e_0, u_0)^\alpha) \quad (12)$$

vanishes identically for every pair of $p+1$ -tuples $(e, u) = (e_0, \dots, e_p, u_0, \dots, u_p)$ of points on L_Γ^{p*} . We will show that it also vanishes for every pair of $p+1$ -tuples of points on $\partial_\infty \mathbb{H}^k$. Since $\chi_{\mathbb{B}_{\mathbb{H}^k}^\alpha |_{L_\Gamma}}^{p_0}$ does not vanish this will finish the proof.

Remark that fixing three points in $\partial_\infty \mathbb{H}^k$ and considering $\mathbb{B}_{\mathbb{H}^k}^\alpha$ as a function on the remaining variable one obtains a rational function on $\partial_\infty \mathbb{H}^k$ to the power α . Consider then common denominator in equation (12) and denote by $Q(e, u)$ the numerator. Remark that $Q(e, u)$ is a sum and subtraction of polynomial functions, each polynomial to a power α .

Fix $\bar{e} = (e_0, \dots, e_p)$ p points in L_Γ and (u_0, \dots, u_{p+1}) $p+1$ points in L_Γ . Consider now the function $f(x) = Q((x, \bar{e}), u) : \partial_\infty \mathbb{H}^k \rightarrow \mathbb{R}$. The function f vanishes identically on L_Γ , hence Proposition 10.5 states that $f \equiv 0$. Repeating this procedure one can pull out one by one the points on L_Γ to $\partial_\infty \mathbb{H}^k$ to show that $\chi_{\mathbb{B}_{\mathbb{H}^k}^\alpha}^p$ vanishes identically on $(\partial_\infty \mathbb{H}^k)^{p*}$. This finishes the proof.

11. PROOFS OF THEOREM B, COROLLARY 1.5 AND COROLLARY 1.6

We will first prove rigidity for the spectral entropy; Hilbert entropy rigidity and Corollary 1.6 follow similarly with minor arrangements. Throughout this section Γ is a Zariski dense convex co-compact group of \mathbb{H}^k , $\rho : \Gamma \rightarrow \mathrm{PGL}(d, \mathbb{R})$ is an irreducible convex representation and $\alpha = \alpha_\rho$ is the best Hölder exponent of the equivariant map $\xi : L_\Gamma \rightarrow \mathbb{P}(\mathbb{R}^d)$.

11.1. Proof of Spectral entropy rigidity. Assume the following equality holds

$$\alpha h_\rho = h_\Gamma \quad (13)$$

and assume that $\mathbf{b}_\rho \geq 0$. Remark 5.2 states that there exists $c > 0$ such that for every non torsion $\gamma \in \Gamma$ one has

$$\lambda_1(\rho\gamma) = c|\gamma|.$$

This equality together with equality (13) imply in fact that

$$\lambda_1(\rho\gamma) = \alpha|\gamma|. \quad (14)$$

The cocycles β_ρ and $\alpha \cdot \text{Busseman} : \Gamma \times L_\Gamma \rightarrow \mathbb{R}$ are hence cohomologous, and thus Lemma 7.3 implies their cross ratios coincide, i.e.: $\mathbf{b}_\rho = |\mathbf{b}_\rho| = \mathbb{B}_{\mathbb{H}^k}^\alpha |L_\Gamma|$.

Since ρ is irreducible, the cross ratio \mathbf{b}_ρ has rank d . Hence proposition 10.3 implies that the cross ratio $\mathbb{B}_{\mathbb{H}^k}^\alpha$, defined on all $\partial_\infty \mathbb{H}^k$, also has rank d . Since $\mathbb{B}_{\mathbb{H}^k}^\alpha$ is invariant under the full isometry group $\text{Isom } \mathbb{H}^k$, Labourie's Theorem 6.6 applies and uniqueness implies that $\rho : \Gamma \rightarrow \text{PGL}(d, \mathbb{R})$ extends to $\bar{\rho} : \text{Isom } \mathbb{H}^k \rightarrow \text{PGL}(d, \mathbb{R})$.

The equivariant map $\xi : L_\Gamma \rightarrow \mathbb{P}(\mathbb{R}^d)$ then comes from a proximal irreducible morphism $\bar{\rho} : \text{Isom } \mathbb{H}^k \rightarrow \text{PGL}(d, \mathbb{R})$ and is thus of class C^∞ , in particular it is Lipschitz i.e. $\alpha = 1$. Hence $\mathbf{b}_{\bar{\rho}} = \mathbb{B}_{\mathbb{H}^k}$, therefore $d = \text{rank } \mathbf{b}_{\bar{\rho}} = \text{rank } \mathbb{B}_{\mathbb{H}^k} = k + 1$. Again Labourie's Theorem 6.6 implies that $\bar{\rho}$ is the Klein model of \mathbb{H}^k .

11.2. Hilbert entropy rigidity. Suppose now that the following equality holds

$$\alpha H_\rho = h_\Gamma.$$

Similar reasoning to spectral entropy rigidity yields

$$\lambda_1(\rho\gamma) - \lambda_d(\rho\gamma) = 2\alpha|\gamma|$$

for every non torsion $\gamma \in \Gamma$. Consider the irreducible adjoint representation $\mathbf{A}_\rho : \Gamma \rightarrow \text{PGL}(V_\rho)$ (recall Lemma 4.7). Again, analogue reasoning to the subsection 11.1 implies that $|\mathbf{b}_{\mathbf{A}_\rho}| = \mathbb{B}_{\mathbb{H}^k}^{2\alpha} |L_\Gamma|$.

Since $\mathbf{b}_\rho \geq 0$ Lemma 6.7 implies that so is $\mathbf{b}_{\mathbf{A}_\rho}$, hence

$$\mathbf{b}_{\mathbf{A}_\rho} = \mathbb{B}_{\mathbb{H}^k}^{2\alpha} |L_\Gamma|.$$

As before, the cross ratio $\mathbb{B}_{\mathbb{H}^k}^{2\alpha}$ has finite rank and thus $\mathbf{A}_\rho : \Gamma \rightarrow \text{PGL}(V_\rho)$ extends to a representation $\bar{\mathbf{A}}_\rho : \text{Isom } \mathbb{H}^k \rightarrow \text{PGL}(V_\rho)$. Hence $\alpha = 1$ and highest restricted weight $\chi_{\bar{\mathbf{A}}_\rho}$ of $\bar{\mathbf{A}}_\rho$ is

$$\chi_{\bar{\mathbf{A}}_\rho}(\lambda_{\mathbb{H}^k}(\gamma)) = 2|\gamma|.$$

Since $\bar{\mathbf{A}}_\rho$ is proximal and irreducible the last equations implies (using Remark 9.3) that $\bar{\mathbf{A}}_\rho$ is the adjoint representation of $\text{PSO}(k, 1)$,

$$\bar{\mathbf{A}}_\rho = \text{Ad} : \text{PSO}(1, k) \rightarrow \text{PGL}(\mathfrak{so}(1, k)).$$

This finishes the proof.

11.3. Proof of Corollary 1.6. Assume now that Γ acts co-compactly on \mathbb{H}^2 and that the equality

$$\alpha h_\rho = 1 \quad (15)$$

holds.

Similarly to Subsection 11.1, $\lambda_1(\rho\gamma) = \alpha|\gamma|$, for every non torsion $\gamma \in \Gamma$, and the cross ratios $|\mathbf{b}_\rho|$ and $\mathbb{B}_{\mathbb{H}^2}^\alpha$ coincide. Hence $|\mathbf{b}_\rho|$ is invariant under group $\mathrm{PSL}(2, \mathbb{R})$. The same happens then for the cross ratio \mathbf{b}_ρ (see Lemma 6.2). Labourie's Theorem 6.6 applies and uniqueness implies that $\rho : \Gamma \rightarrow \mathrm{PGL}(d, \mathbb{R})$ extends to $\bar{\rho} : \mathrm{PSL}(2, \mathbb{R}) \rightarrow \mathrm{PGL}(d, \mathbb{R})$. Hence α equals 1.

Let χ_ρ be the restricted highest weight of the irreducible representation $\bar{\rho} : \mathrm{PSL}(2, \mathbb{R}) \rightarrow \mathrm{PGL}(d, \mathbb{R})$. Equation (11) states that

$$\chi_\rho(\lambda_{\mathbb{H}^2}(\gamma)) = \lambda_1(\rho\gamma) = |\gamma|.$$

Proposition 9.1 implies that $\bar{\rho}$ is the Klein model of \mathbb{H}^2 . This finishes the proof.

12. HYPERCONVEX REPRESENTATIONS AND THEOREM C

We will freely use the notations of Section 9. Let G be a real non compact semi-simple Lie group and denote by \mathcal{F} the *Furstenberg boundary* of the symmetric space of G . The product $\mathcal{F} \times \mathcal{F}$ has a unique open G -orbit denoted by $\mathcal{F}^{(2)}$.

Definition 12.1. A representation $\rho : \Gamma \rightarrow G$ is *hyperconvex* if there exists a ρ -equivariant Hölder-continuous map $\zeta : \partial_\infty \Gamma \rightarrow \mathcal{F}$ such that if $x \neq y$ are distinct points in $\partial_\infty \Gamma$ then the pair $(\zeta(x), \zeta(y))$ belongs to $\mathcal{F}^{(2)}$.

The following Lemma relates hyperconvex representations to convex ones.

Lemma 12.2. *If $\rho : \Gamma \rightarrow G$ is Zariski dense and hyperconvex, and $\Lambda : G \rightarrow \mathrm{PGL}(V)$ is a finite dimensional irreducible proximal representation, then the composition $\Lambda \circ \rho : \Gamma \rightarrow \mathrm{PGL}(V)$ is irreducible and convex.*

Proof. A proximal representation $\Lambda : G \rightarrow \mathrm{PGL}(V)$ induces a C^∞ equivariant map $\mathcal{F} \rightarrow \mathbb{P}(V)$. Considering the dual representation $\Lambda^* : G \rightarrow \mathrm{PGL}(V^*)$ one obtains another equivariant map $\mathcal{F} \rightarrow \mathrm{PGL}(V^*)$. The remainder of the statement is direct. \square

We need the following Theorem from [20].

Theorem 12.3 ([20, Corollary 7.6]). *Let $\rho : \Gamma \rightarrow G$ be a Zariski dense hyperconvex representation, then there exists a Γ -invariant Hölder-continuous function $F : \mathrm{U}\Gamma \rightarrow \mathfrak{a}$ such that for every non torsion conjugacy class $[\gamma] \in [\Gamma]$ one has*

$$\int_{[\gamma]} F = \lambda(\rho\gamma).$$

Moreover, if $\varphi|\mathfrak{a}^+ - \{0\} > 0$ then $\varphi(F) : \mathrm{U}\Gamma \rightarrow \mathbb{R}$ is Livšic-cohomologous to a positive function.

Fix an action of Γ on $\mathrm{CAT}(-1)$ space X , and denote L_Γ its limit set on $\partial_\infty X$. Assume from now on that $\rho : \Gamma \rightarrow G$ is a Zariski dense hyperconvex representation and denote by α the Hölder exponent of the equivariant map $\zeta : L_\Gamma \rightarrow \mathcal{F}$.

Lemma 12.4. *For every simple root $\theta \in \Pi$ and every non torsion $\gamma \in \Gamma$ one has*

$$\alpha \leq \frac{\theta(\lambda(\rho\gamma))}{|\gamma|}.$$

Proof. Consider $\Lambda_\theta \circ \rho : \Gamma \rightarrow \mathrm{PGL}(V_\theta)$ the irreducible convex representation given by Tits's Proposition 9.2 and Lemma 12.2. One then has

$$\theta(\lambda(\rho\gamma)) = \lambda_1(\Lambda_\theta \circ \rho\gamma) - \lambda_2(\Lambda_\theta \circ \rho\gamma).$$

The Lemma follows from Lemma 4.11. \square

12.1. Proof of Theorem C. The proof is very similar to the proof of Theorem A. Consider $F : \mathrm{U}\Gamma \rightarrow \mathfrak{a}$ given by Theorem 12.3, i.e. F verifies

$$\int_{[\gamma]} F = \lambda(\rho\gamma)$$

for every non torsion conjugacy class $[\gamma]$ of Γ .

Let φ be a linear functional such that $\varphi|_{\mathfrak{a}^+ - \{0\}} > 0$. Using Theorem 12.3 we may assume that $f = \varphi(F) : \mathrm{U}\Gamma \rightarrow \mathbb{R}_+^*$ a positive function. Remark that

$$\varphi(\lambda(\rho\gamma)) = \int_{[\gamma]} f$$

and hence $h_f = h_\varphi$. Consider a sequence of conjugacy classes $[\gamma_n]$ such that

$$\frac{\mathrm{Leb}_{\gamma_n}}{|\gamma_n|} \rightarrow m_{-h_\varphi f}$$

as $n \rightarrow \infty$, where $m_{-h_\varphi f}$ denotes the equilibrium state of $-h_\varphi f$. Analog reasoning to Theorem A shows that

$$\frac{\varphi(\lambda(\rho\gamma_n))}{|\gamma_n|} \leq \frac{h(\phi, m_{-h_\varphi f})}{h_\varphi} (1 + \varepsilon),$$

and thus, using Lemma 12.4 one finds that

$$\frac{\alpha h_\varphi}{h_\Gamma} \leq \frac{\alpha h_\varphi}{h(\phi, m_{-h_\varphi f})} \leq \frac{\theta(\lambda(\rho\gamma_n))}{\varphi(\lambda(\rho\gamma_n))} (1 + \varepsilon)$$

for every simple root $\theta \in \Pi$. We now try to maximize the function $V : \mathbb{P}(\mathfrak{a}^+) \rightarrow \mathbb{R}$

$$V(a) = \min_{\theta \in \Pi} \left\{ \frac{\theta(a)}{\varphi(a)} \right\}.$$

We need the following Linear Algebra Lemma. Consider an n -dimensional vector space W , a k -simplex is the convex hull of $k+1$ points $\{x_0, \dots, x_k\}$ in W such that for every $i \in \{0, \dots, k\}$ the set $\{x_0, \dots, x_k\} - \{x_i\}$ is linearly independent.

Lemma 12.5. *Consider $n+1$ affine linear functionals $\varphi_i : W \rightarrow \mathbb{R}$ on an n -dimensional vector space V , such that*

$$\Delta := \bigcap_0^n \{v \in W : \varphi_i(v) \geq 0\}$$

is an n -dimensional simplex. Then

$$\max_{v \in \Delta} \min\{\varphi_i(v) : i \in \{0, \dots, n\}\}$$

is given in the point all the φ_i 's coincide, i.e. in the unique $v \in \Delta$ such that

$$\varphi_0(v) = \varphi_1(v) = \dots = \varphi_n(v).$$

Proof. The proof of the Lemma is trivial. \square

Fix now some vector v in the interior of \mathfrak{a}^+ such that $\varphi(v) \neq 0$ and consider the map $T : \ker \varphi \rightarrow \mathbb{P}(\mathfrak{a})$ given by $w \mapsto \mathbb{R}(v + w)$, this map identifies $\ker \varphi$ with $\mathbb{P}(\mathfrak{a}) - \mathbb{P}(\ker \varphi)$. Via this mapping one gets that the functions $T_\theta : \ker \varphi \rightarrow \mathbb{R}$ given by

$$T_\theta(w) := \frac{\theta(w + v)}{\varphi(w + v)} = \frac{\theta(v)}{\varphi(v)} + \frac{\theta(w)}{\varphi(v)}$$

are affine functionals. Since φ is positive on the Weyl chamber $\mathfrak{a}^+ - \{0\}$ we get that

$$\Delta = T^{-1}(\mathbb{P}(\mathfrak{a}^+)) = T^{-1}(\mathbb{P}(\bigcap_{\theta \in \Pi} \{\theta \geq 0\})) = \bigcap_{\theta \in \Pi} \{T_\theta \geq 0\}$$

is a simplex of dimension $\dim \mathfrak{a} - 1 = \dim \ker \varphi$.

Remark that $V \circ T = \min\{T_\theta : \theta \in \Pi\}$, hence Lemma 12.5 implies that the maximum of $V \circ T|_\Delta$ is realized where all the functions $\{T_\theta : \theta \in \Pi\}$ coincide, i.e. in the set

$$\{a \in \mathfrak{a}^+ : \theta_1(a) = \theta_2(a) \text{ for every pair } \theta_1, \theta_2 \in \Pi\}.$$

This is exactly the barycenter of the Weyl chamber $\text{bar}_{\mathfrak{a}^+}$.

Hence

$$\frac{\alpha h_\varphi}{h_\Gamma} \leq \frac{\alpha h_\varphi}{h(\phi, m_{-h_\varphi f})} \leq V(\lambda(\rho\gamma_n))(1 + \varepsilon) \leq \frac{\theta(\text{bar}_{\mathfrak{a}^+})}{\varphi(\text{bar}_{\mathfrak{a}^+})}(1 + \varepsilon). \quad (16)$$

This shows the desired inequality.

Remark 12.6. As in Theorem A, Remark that equality in equation (16) implies that f is Livšic cohomologous to a constant.

13. PROOFS OF THEOREM D AND COROLLARY 1.12

Assume now that Γ is a Zariski dense convex-co-compact group of \mathbb{H}^k . Consider a hyperconvex representation $\rho : \Gamma \rightarrow G$ and consider χ , a restricted weight of G . Suppose the equality

$$\alpha_\rho h_\chi = h_\Gamma \frac{\theta(\text{bar}_{\mathfrak{a}^+})}{\chi(\text{bar}_{\mathfrak{a}^+})}$$

holds, where θ is any simple root.

Remark 12.6 implies that the function $f_\chi = \chi(F) : \text{U}\Gamma \rightarrow \mathbb{R}_+^*$ such that

$$\chi(\lambda(\rho\gamma)) = \int_{[\gamma]} f_\chi$$

is Livšic cohomologous to a constant. Consider now the proximal irreducible representation $\Lambda_\chi : G \rightarrow \text{PGL}(V_\chi)$ associated to χ by Proposition 9.1, and the composition

$$\phi = \Lambda_\chi \circ \rho : \Gamma \rightarrow \text{PGL}(V_\chi).$$

Lemma 12.2 states that ϕ is irreducible and convex and by definition one has

$$\lambda_1(\phi\gamma) = \chi(\lambda(\rho\gamma)) = c|\gamma|.$$

The proof now follows similar to Theorem B. Consider \mathbf{b}_ϕ the cross ratio on $L_\Gamma \subset \partial_\infty \mathbb{H}^k$ of the representation ϕ . The last equality together with Lemma 7.3 imply that $|\mathbf{b}_\phi| = \mathbb{B}_{\mathbb{H}^k}^c|L_\Gamma$.

We now distinguish the two situations:

- Assume $\mathbf{b}_\phi \geq 0$ (i.e. we are in the hypothesis of Theorem D): We then have that $\mathbf{b}_\phi = \mathbb{B}_{\mathbb{H}^k}^c|_{L_\Gamma}$. Since ϕ is irreducible, the cross ratio \mathbf{b}_ϕ has finite rank and Proposition 10.3 implies that the cross ratio $\mathbb{B}_{\mathbb{H}^k}^c$, defined on $\partial_\infty \mathbb{H}^k$, has the same rank. Hence Labourie's Theorem 6.6 applies and uniqueness implies that $\phi : \Gamma \rightarrow \mathrm{PGL}(V_\chi)$ extends to $\bar{\phi} : \mathrm{Isom} \mathbb{H}^k \rightarrow \mathrm{PGL}(V_\chi)$.
- Γ is co-compact on \mathbb{H}^2 (i.e. we are in the hypothesis of Corollary 1.12): In this case we have $|\mathbf{b}_\phi| = \mathbb{B}_{\mathbb{H}^2}^c$, and thus $|\mathbf{b}_\phi|$ is invariant under the group $\mathrm{PSL}(2, \mathbb{R})$. The same happens then for the cross ratio \mathbf{b}_ϕ (see Lemma 6.2). Labourie's Theorem 6.6 applies and uniqueness implies that $\phi : \Gamma \rightarrow \mathrm{PGL}(V_\chi)$ extends to $\bar{\phi} : \mathrm{PSL}(2, \mathbb{R}) \rightarrow \mathrm{PGL}(V_\chi)$.

In both cases we get that $\phi : \Gamma \rightarrow \mathrm{PGL}(V_\chi)$ extends to $\bar{\phi} : \mathrm{Isom} \mathbb{H}^k \rightarrow \mathrm{PGL}(V_\chi)$. The Zariski closure of $\phi(\Gamma)$ is $\bar{\phi}(\mathrm{Isom} \mathbb{H}^k)$ on one hand, and $\Lambda_\chi(G)$ on the other. Hence $\Lambda_\chi(G) = \bar{\phi}(\mathrm{Isom} \mathbb{H}^k)$. This finishes the proof.

14. PROOF OF COROLLARY 1.13

We will now prove the following Corollary.

Corollary. *Let $\rho : \pi_1 \Sigma \rightarrow \mathrm{PSL}(d, \mathbb{R})$ be a representation in the Hitchin component and denote by α the best Hölder exponent of the equivariant map $\zeta : \partial_\infty \mathbb{H}^2 \rightarrow \mathcal{F}$, then*

$$\alpha h_\rho \leq \frac{2}{d-1} \text{ and } \alpha_\rho H_\rho \leq \frac{2}{d-1}$$

and either equality holds only if $\rho = \tau_d \circ \mathbf{f}$, where $\tau_d : \mathrm{PSL}(2, \mathbb{R}) \rightarrow \mathrm{PSL}(d, \mathbb{R})$ is the irreducible representation and $\mathbf{f} : \pi_1 \Sigma \rightarrow \mathrm{PSL}(2, \mathbb{R})$ is the departing action.

The proof goes as follows. Denote by G the Zariski closure of ρ . Guichard's Theorem 14.1 below implies that G is simple, hence $\rho : \pi_1 \Sigma \rightarrow G$ is again hyperconvex. Consider \mathfrak{a} a Cartan subspace of \mathfrak{g} , and let $\chi \in \mathfrak{a}^*$ be the restricted highest weight of the (irreducible proximal) representation $G \subset \mathrm{PSL}(d, \mathbb{R})$, i.e. if $g \in G$ then $\chi(\lambda(g)) = \lambda_1(g)$.

Remark that by definition the entropy of ρ relative to χ is the spectral entropy $h_\rho = h_\chi$ of ρ , and the entropy of ρ relative to

$$\varphi = \frac{\chi + \chi \circ \mathbf{i}}{2}$$

is the Hilbert entropy $H_\rho = h_\varphi$ of ρ . We will prove the Corollary for the spectral entropy, the other being completely analogous.

Theorem C asserts that

$$\alpha h_\rho \leq \frac{\theta(\mathrm{bar}_{\mathfrak{a}^+})}{\chi(\mathrm{bar}_{\mathfrak{a}^+})} \tag{17}$$

for any simple root $\theta \in \Pi$ of \mathfrak{a} and where $\mathrm{bar}_{\mathfrak{a}^+}$ is the barycenter of the Weyl chamber \mathfrak{a}^+ . Theorem D implies that equality in (17) can only hold if G is isomorphic to $\mathrm{PSL}(2, \mathbb{R})$.

Guichard's Theorem gives a finite list of possible groups G , i.e. of possible Zariski closures of $\rho(\pi_1 \Sigma)$. We will finish by an explicit computation showing that in all possible cases one has

$$\frac{\theta(\mathrm{bar}_{\mathfrak{a}^+})}{\chi(\mathrm{bar}_{\mathfrak{a}^+})} = \frac{2}{d-1}.$$

The author would like to thank Olivier Guichard for discussions concerning his work.

Theorem 14.1 (Guichard [13]). *Let $\rho : \pi_1 \Sigma \rightarrow \mathrm{SL}(d, \mathbb{R})$ be the lift of a representation in the Hitchin component, then the Zariski closure $\overline{\rho^\mathbb{Z}}$ is either conjugate to $\tau_d(\mathrm{SL}(2, \mathbb{R}))$, $\mathrm{SL}(d, \mathbb{R})$ or conjugate to one of the following groups:*

- $\mathrm{Sp}(2n, \mathbb{R})$ if $d = 2n$,
- $\mathrm{SO}(n, n+1)$ if $d = 2n+1$,
- G_2 or $\mathrm{SO}(3, 4)$ if $d = 7$.

For $i \in \{1, \dots, k\}$ we will denote by $\varepsilon_i : \mathbb{R}^k \rightarrow \mathbb{R}$ the function

$$\varepsilon_i(a_1, \dots, a_k) = a_i.$$

We refer the reader to Knapp's book [16] for the standard computations of simple roots and highest weights that follow.

The $\tau_d(\mathrm{SL}(2, \mathbb{R}))$ and $\mathrm{SL}(d, \mathbb{R})$ cases. Assume first that $\rho(\pi_1 \Sigma)$ is Fuchsian, i.e. it is Zariski dense in $\tau_d(\mathrm{SL}(2, \mathbb{R}))$. A Cartan subspace of $\mathfrak{sl}(2, \mathbb{R})$ is $\mathfrak{a} = \{(a, -a) : a \in \mathbb{R}\}$ the Weyl chamber is $\mathfrak{a}^+ = \{(a, -a) : a \geq 0\}$ with simple root $\Pi = \{2\varepsilon_1\}$. The highest weight of the representation τ_d is $\chi(a, -a) = (d-1)a$. Hence

$$\frac{\theta(\mathrm{bar}_{\mathfrak{a}^+})}{\chi(\mathrm{bar}_{\mathfrak{a}^+})} = \frac{2a}{(d-1)a} = \frac{2}{d-1}.$$

Suppose now that $\rho(\pi_1 \Sigma)$ is Zariski dense in $\mathrm{SL}(d, \mathbb{R})$. The Cartan subspace of $\mathfrak{sl}(d, \mathbb{R})$ is $\mathfrak{a} = \{(a_1, \dots, a_d) \in \mathbb{R}^d : a_1 + \dots + a_d = 0\}$ and

$$\mathfrak{a}^+ = \{(a_1, \dots, a_d) \in \mathfrak{a} : a_1 \geq \dots \geq a_d\},$$

the simple roots are

$$\Pi = \{\theta_i(a_1, \dots, a_d) = a_i - a_{i+1} : i \in \{1, \dots, d-1\}\}$$

and the barycenter is

$$\mathrm{bar}_{\mathfrak{a}^+} = \{((d-1)t, (d-3)t, \dots, (3-d)t, (1-d)t) : t \geq 0\}.$$

Hence for any $\theta \in \Pi$ one has

$$\frac{\theta(\mathrm{bar}_{\mathfrak{a}^+})}{\chi(\mathrm{bar}_{\mathfrak{a}^+})} = \frac{2t}{(d-1)t} = \frac{2}{d-1}.$$

The $\mathrm{Sp}(2n, \mathbb{R})$ case. Assume $d = 2n$ and that the Zariski closure of $\rho(\pi_1 \Sigma)$ is $\mathrm{Sp}(2n, \mathbb{R})$. Standard computations show that $\mathfrak{a} = \mathbb{R}^n$, and a Weyl chamber is

$$\mathfrak{a}^+ = \{(a_1, \dots, a_n) : a_i \geq a_{i+1} \text{ } i = 1, \dots, n-1 \text{ and } a_n \geq 0\}.$$

The set of simple roots associated to this Weyl chamber is

$$\Pi = \{\varepsilon_i - \varepsilon_{i+1} : i = 1, \dots, n-1\} \cup \{2\varepsilon_n\}.$$

The barycenter of the Weyl chamber is hence

$$\mathrm{bar}_{\mathfrak{a}^+} = \{((2n-1)t, (2n-3)t, \dots, 3t, t) : t \geq 0\}.$$

The highest weight of the representation $\mathrm{Sp}(2n, \mathbb{R}) \subset \mathrm{SL}(d, \mathbb{R})$ is $\chi(a_1, \dots, a_n) = a_1$. Finally, for any $\theta \in \Pi$ one has

$$\frac{\theta(\mathrm{bar}_{\mathfrak{a}^+})}{\chi(\mathrm{bar}_{\mathfrak{a}^+})} = \frac{2t}{(2n-1)t} = \frac{2}{d-1}.$$

The $\mathrm{SO}(n, n+1)$ case. Suppose now that $d = 2n + 1$ and that the Zariski closure of $\rho(\pi_1 \Sigma)$ is $\mathrm{SO}(n, n+1)$. Standard computations show that $\mathfrak{a} = \mathbb{R}^n$, and a Weyl chamber is

$$\mathfrak{a}^+ = \{(a_1, \dots, a_n) : a_i \geq a_{i+1} \text{ } i = 1, \dots, n-1 \text{ and } a_n \geq 0\}.$$

The set of simple roots associated to this Weyl chamber is

$$\Pi = \{\varepsilon_i - \varepsilon_{i+1} : i = 1, \dots, n-1\} \cup \{\varepsilon_n\}.$$

The barycenter of the Weyl chamber is hence

$$\mathrm{bar}_{\mathfrak{a}^+} = \{(nt, (n-1)t, \dots, 2t, t) : t \geq 0\}.$$

The highest weight of the representation $\mathrm{SO}(n, n+1) \subset \mathrm{SL}(d, \mathbb{R})$ is $\chi(a_1, \dots, a_n) = a_1$. Finally, for any $\theta \in \Pi$ one has

$$\frac{\theta(\mathrm{bar}_{\mathfrak{a}^+})}{\chi(\mathrm{bar}_{\mathfrak{a}^+})} = \frac{t}{nt} = \frac{1}{n} = \frac{2}{d-1}.$$

The G_2 case. The remaining case is $d = 7$ and the Zariski closure of $\rho(\pi_1 \Sigma)$ being the exceptional simple Lie group G_2 . We refer the reader to Knapp's book [16, page 692] for the following computations. In this case we have

$$\mathfrak{a} = \{(a_1, a_2, a_3) \in \mathbb{R}^3 : a_1 + a_2 + a_3 = 0\},$$

a Weyl chamber is

$$\mathfrak{a}^+ = \{(a_1, a_2, a_3) : a_1 \geq a_2 \text{ and } -2a_1 + a_2 + a_3 \geq 0\}.$$

The set of simple roots is

$$\Pi = \{\varepsilon_1 - \varepsilon_2, -2\varepsilon_1 + \varepsilon_2 + \varepsilon_3\},$$

and the barycenter of the Weyl chamber is hence

$$\mathrm{bar}_{\mathfrak{a}^+} = \{(-t, -4t, 5t) : t \geq 0\}.$$

The highest weight associated to the representation $\mathrm{G}_2 \rightarrow \mathrm{SL}(7, \mathbb{R})$ is

$$\chi = \omega_1 = 2(\varepsilon_1 - \varepsilon_2) - 2\varepsilon_1 + \varepsilon_2 + \varepsilon_3 = \varepsilon_3 - \varepsilon_2.$$

Finally, for any $\theta \in \Pi$ one has

$$\frac{\theta(\mathrm{bar}_{\mathfrak{a}^+})}{\chi(\mathrm{bar}_{\mathfrak{a}^+})} = \frac{3t}{5t+4t} = \frac{1}{3} = \frac{2}{d-1}.$$

This finishes the proof.

REFERENCES

- [1] L.M. Abramov. On the entropy of a flow. *Dokl. Akad. Nauk. SSSR*, 128:873–875, 1959.
- [2] Y. Benoist. Propriétés asymptotiques des groupes linéaires. *Geom. funct. anal.*, 7(1):1–47, 1997.
- [3] Y. Benoist. Propriétés asymptotiques des groupes linéaires ii. *Adv. Stud. Pute Math.*, 26:33–48, 2000.
- [4] Y. Benoist. Convexes divisibles I. In *Algebraic groups and arithmetic*, pages 339–374. Tata Inst. Fund. Res., 2004.
- [5] Y. Benoist. Convexes divisibles III. *Ann. Sci. École Norm. Sup.*, 38:793–832, 2005.
- [6] M. Bourdon. Structure conforme au bord et flot géodésique d'un $\mathrm{CAT}(-1)$ -espace. *Enseign. Math. (2)*, 41:63–102, 1995.
- [7] M. Bourdon. Sur le birapport au bord des $\mathrm{CAT}(-1)$ spaces. *Publ. Math. I.H.E.S.*, 83:95–104, 1996.

- [8] B. H. Bowditch. Cut points and canonical splittings of hyperbolic groups. *Acta Mathematica*, 180(2):145–186, 1998.
- [9] R. Bowen and D. Ruelle. The ergodic theory of axiom A flows. *Invent. Math.*, 29:181–202, 1975.
- [10] M. Bridgeman, D. Canary, F. Labourie, and A. Sambarino. The pressure metric for convex representations. preprint.
- [11] G. Courtois. Critical exponents and rigidity in negative curvature. In *Séminaires & Congrès, Géométries à courbure négative ou nulle, groupes discrets et rigidité*, volume 18, pages 293–319. Société mathématique de France, 2009.
- [12] M. Crampon. Entropies of compact strictly convex projective manifolds. *J. of Mod. Dyn.*, 3:511–547, 2009.
- [13] O. Guichard. Personal communication.
- [14] O. Guichard and A. Weinhard. Anosov representations: domains of discontinuity and applications. *Invent. Math.*, 190:357–438, 2012. ArXiv: 1108.0733v1.
- [15] N. J. Hitchin. Lie groups and Teichmüller space. *Topology*, 31(3):449–473, 1992.
- [16] A. Knapp. *Lie groups beyond an introduction*. Birkhäuser, 2002.
- [17] F. Labourie. Anosov flows, surface groups and curves in projective space. *Invent. Math.*, 165:51–114, 2006.
- [18] F. Labourie. Cross ratios, surface groups, $SL(n, \mathbb{R})$ and diffeomorphisms of the circle. *Publ. Math. I.H.E.S.*, 106:139–213, 2007.
- [19] F. Ledrappier. Structure au bord des variétés à courbure négative. *Séminaire de théorie spectrale et géométrie de Grenoble*, 71:97–122, 1994-1995.
- [20] A. Sambarino. Quantitative properties of convex representations. To appear in *Comment. Math. Helv.* arXiv:1104.4705v1, 2011.
- [21] M. Shub. *Global stability of dynamical systems*. Springer Verlag, New York, 1987.
- [22] D. Sullivan. The density at infinity of a discrete group of hyperbolic motions. *Publ. Math. de l’I.H.E.S.*, 50:171–202, 1979.
- [23] J. Tits. Représentations linéaires irréductibles d’un groupe réductif sur un corps quelconque. *J. Reine Angew. Math.*, 247:196–220, 1971.
- [24] C. Yue. The ergodic theory of discrete isometry groups on manifolds of variable negative curvature. *Trans. of the A.M.S.*, 348(12):4965–5005, 1996.

Andrés Sambarino
 Departement de Mathématiques
 Université Paris Sud,
 F-91405 Orsay France,
 andres.sambarino@gmail.com